

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
5 April 2001 (05.04.2001)

PCT

(10) International Publication Number
WO 01/23411 A2

- (51) International Patent Classification⁷: C07K [US/US]; 31 Mayflower Drive, Wenham, MA 01984 (US).
GARDNER, Andrew, F. [US/US]; 1163 Beacon Street, Apartment #6, Brookline, MA 02446 (US).
- (21) International Application Number: PCT/US00/26900
- (22) International Filing Date: 29 September 2000 (29.09.2000) (74) Agent: WILLIAMS, Gregory, D.; New England Biolabs, Inc., 32 Tozer Road, Beverly, MA 01915 (US).
- (25) Filing Language: English (81) Designated States (*national*): JP, US.
- (26) Publication Language: English (84) Designated States (*regional*): European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).
- (30) Priority Data: 60/157,204 30 September 1999 (30.09.1999) US
- (71) Applicant (*for all designated States except US*): NEW ENGLAND BIOLABS, INC. [US/US]; 32 Tozer Road, Beverly, MA 01915 (US).
- Published:
— Without international search report and to be republished upon receipt of that report.
- (72) Inventors; and
(75) Inventors/Applicants (*for US only*): JACK, William, E.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: INCORPORATION OF MODIFIED NUCLEOTIDES BY ARCHAEON DNA POLYMERASES AND RELATED METHODS

(57) Abstract: The present invention is directed toward improving the efficiency of chain terminator incorporation by Family B archaeon DNA polymerases. Previously, the low efficiency of ddNTP, and more especially dye-labeled ddNTP, incorporation has limited the usefulness of this group of DNA polymerases in protocols requiring chain terminator incorporation.

WO 01/23411 A2

INCORPORATION OF MODIFIED NUCLEOTIDES BY ARCHAEON DNA POLYMERASES AND RELATED METHODS

BACKGROUND OF THE INVENTION

5

DNA polymerases have played a central role in the development of molecular biology. Their use is at the core of a wide range of laboratory protocols, including DNA sequencing (Sanger, *et al.*, *Proc. Natl. Acad. Sci., USA* **74**:5463-5467 (1977)), strand displacement amplification (SDA; Walker, *et al.*, *Proc. Natl. Acad. Sci., USA* **89**:392-396 (1992)), probe labeling, site-directed mutagenesis, the polymerase chain reaction (PCR; Saiki, *et al.*, *Science*, **230**:1350-1354 (1985)), and cloning. These applications depend critically on the ability of polymerases to faithfully replicate DNA, either to create a product whose biological properties are identical to the substrate, or to create a product whose identity accurately reflects the substrate, thus facilitating characterization and manipulation of this substrate.

10

15

20

25

30

35

A number of applications require polymerases that are able to incorporate modified nucleotides. One such application is chain terminator nucleic acid sequencing where nucleotides with modified sugars, most often a dideoxynucleotide (ddNTP), are employed to deduce the ordering of bases in a sequencing sample (Sanger, *et al.*, *supra.* (1977)). Sequence-specific chain termination, occurring upon incorporation of these analogs, creates a product whose length measures the position of the complementary base in the substrate molecule. By ordering such products derived using terminators corresponding to A, G, C and T bases, the nucleic acid sequence can be deduced. Production of these terminator products can occur in four parallel reactions, each with a single A, G, C or T terminator, to deduce the DNA sequence. Alternatively, if each of the four terminators contains a unique detection agent, the terminator products can be produced simultaneously in one reaction. One family of such detection agents is fluorescent probes (Prober, *et al.*, *Science* **238**:336-341 (1987); US Patent No. 5,332,666). Fluorescent probes can either be used individually or, if the emission spectra are distinguishable, in multiple sets. Fluorescent probes are most often

-2-

attached to the nucleotide base, creating a need in the art for DNA polymerases that can readily incorporate such dye-labeled nucleotides. Detection probes can also be moieties that interact with a second molecule, such as an antibody, with indirect detection occurring via the second molecule. Such is the case, for example, with the binding of specific antibodies to fluorescein or the binding of streptavidin to biotin. Finally, detection probes can also be radioisotopes, detectable by such methods as autoradiography.

Modifications of chain terminator DNA sequencing have been described that focus on single nucleotide loci, allowing detection of single nucleotide polymorphisms (SNPs; Sarkar, *et al.*, *Genomics* **13**:441-443 (1992); Nikiforov, *et al.*, *Nucleic Acids Res.* **22**:4167-4175; Chen and Kwok, *Nucleic Acids Res.* **15**:347-353 (1997); Chen, *et al.*, *Genome Research* **9**:492-498 (1999); US Patent Nos. 5,888,819, 5,952,174, 6,004,744, and 6,013,43). Such single base detection methods have been instructive in genetic testing and analysis, and also require DNA polymerases able to incorporate modified nucleotides with attached detection probes and/or chain terminating functionalities.

A difficulty with methods requiring the incorporation of modified nucleotides is the inherent fidelity of DNA polymerases. As might be expected, incorporation of a number of nucleotide analogs by DNA polymerases is less efficient than incorporation of the naturally occurring residues, namely dATP, dCTP, dGTP and TTP. As a consequence, technologies relying on incorporation of the analogs can suffer from incomplete and non-uniform incorporation. Accordingly, there is a need in the art for DNA polymerases and nucleotide analog combinations that allow for ready incorporation while retaining base specificity. Since a number of methods require a step in which the DNA is denatured at high temperatures, there is a need for such enzymes that are additionally thermostable.

Several approaches can be envisioned to enhance incorporation of modified nucleotides *in vitro*. First, use of DNA polymerases that more readily incorporate modified nucleotides. Second, use of DNA polymerase variants that more readily incorporate modified nucleotides. Third, use of

-3-

nucleotide analogs that are more readily incorporated by the target DNA polymerase. Fourth, use of high concentrations of modified nucleotides, driving incorporation by the law of mass action. This final approach is limited by the requirement for larger quantities of reagent, and from the higher detection background introduced by the unincorporated nucleotide, and thus is not the preferred approach.

Although in theory any DNA polymerase could be used to incorporate modified nucleotides, polymerases derived from different sources can have different spectra of nucleotide and nucleotide analog incorporation efficiencies. Thus, the choice of polymerase is important in analog incorporation. Primary amino acid sequence similarities allow the classification of most DNA polymerases into three Families, A, B and C, according to similarities with *Escherichia coli* polymerases I, II and III, respectively (Ito and Braithwaite, *Nucleic Acids Res.*, **19**:4045-4057 (1991); Heringa and Argos, *The Evolutionary Biology of Viruses*, Morse, S.S., ed., pp. 87-103, Raven Press, New York (1992)). DNA polymerases of Family A have been the predominant enzymes used in DNA sequence determination, and thus have been most extensively studied with regards to their ability to incorporate chain terminators and dye-labeled nucleotides.

A comparison of two Family A DNA polymerases, the Klenow fragment of DNA polymerase I and T7 DNA polymerase, revealed a remarkable difference in incorporation of chain terminating dideoxynucleoside triphosphate (ddNTPs) (Tabor and Richardson, *Proc. Natl. Acad. Sci., USA* **92**:6339-43 (1995)). Further work established that replacement of F762 in Klenow by tyrosine, the residue present in the analogous position in T7 DNA polymerase, drastically increased the efficiency of ddNTP utilization by that enzyme (Tabor and Richardson, *supra.*). Replacement of the analogous residue of the thermostable Family A Taq DNA polymerase (F667) gave a similar increase in incorporation efficiency of ddNTPs (US Patent No. 5,614,365). These examples illustrate both the use of alternate polymerases and of polymerase variants to increase the efficiency of terminator incorporation.

DNA polymerases of other families could also be considered for incorporation of modified nucleotides. Since a number of applications involve a heat step for DNA strand denaturation, thermostable enzymes of Family B have been explored as candidates for incorporation of modified nucleotides, including a number derived from thermophilic archaea. Such enzymes include, but are not limited to, Vent® DNA polymerase, originally isolated from *Thermococcus litoralis* (Perler, *et al.*, *Proc. Natl. Acad. Sci. USA* **89**:5577-5581 (1992); US Patent Nos. 5,500,363, 5,834,285, 5,352,778); *Pyrococcus furiosus* (*Pfu*) DNA polymerase (US Patent Nos. 5,489,523, 5,827,716), Deep Vent® DNA polymerase (US Patent No. 5,834,285), *Thermococcus barossii* (*Tba*) DNA polymerase (US Patent No. 5,882,904) and 9°N™ DNA polymerase (Southworth, *et al.*, *Proc. Natl. Acad. Sci. USA* **93**: 5281-5285 (1996) and U.S. Patent No. 5,756,334).

Early experiments suggested that archaeon DNA polymerases were not promising candidates for applications requiring incorporation of modified nucleotides such as ddNTPs, more particularly those labeled with dyes. Those enzymes for which kinetic information is available, Vent® and Deep Vent®, both have a relatively high K_m for nucleotides (Kong, *et al.*, *J. Biol. Chem.* **268**:1965-1975 (1993); *New England Biolabs Catalog* 1998/1999, p. 73), and at least Vent® and *Pfu* incorporate unsubstituted ddNTP terminators inefficiently (Gardner and Jack, *Nucleic Acids Res.* **27**:2545-2553 (1999); US Patent No. 5,827,716). Furthermore, reports indicated poor incorporation of dye-substituted ddNTP terminators in DNA sequencing reactions ("CircumVent®: Questions and Answers," *The NEB Transcript*, September 1992, p. 12-13) and in arrayed-primer extension format reactions involving dye-labeled dideoxy terminators (*Tba* DNA polymerase, US Patent No. 5,882,904).

The thermostable archaeon DNA polymerases are not alone in having difficulty incorporating dye-labeled ddNTPs. For example, even though incorporation of ddNTPs is dramatically increased in F667Y versions of *Taq* DNA polymerase (US Patent No. 5,614,365; also know by the trade names Thermo Sequenase™ (Amersham Pharmacia Biotech, Piscataway, NJ) and AmpliTaq® DNA Polymerase, FS (Perkin-Elmer)), dye-terminator

incorporation is still characterized by "...less uniform peak height patterns when compared to primer chemistry profiles, suggesting that the dyes and/or their linker arms affect enzyme selectivity." (Brandis, *Nucleic Acids Res.* 27:1912-1918 (1999)).

Returning to the case of Vent® DNA polymerase, limited information suggests that certain dye-labeled nucleotides can be incorporated. US Patent No. 5,723,298 claims to have used the infrared dye-labeled IRD40 dATP as a substrate for polymerization by CircumVent® thermostable polymerase, although no quantitative aspects of the polymerization were disclosed. CircumVent® is a trade name referring to Vent® DNA polymerase (exo-), a 3'-5' exonuclease-deficient form of Vent® DNA polymerase (New England Biolabs, Beverly, MA). More recently, Ilsey and Buzby presented data at the American Society of Biochemistry and Molecular Biology meeting in May of 1999 regarding incorporation of several dye-substituted nucleotides by a variety of DNA polymerases, including *Taq* and Vent® (exo-) DNA polymerases (Ilsey and Buzby, *The FASEB J.* 13:A1441 (1999)). Incorporation of nine indocyanine and rhodamine dCTP analogs by five polymerases was evaluated. In the case of Vent® (exo-) and *Taq* DNA polymerases, two dye-substituted dCTP analogs were identified whose incorporation was preferred over dCTP: the indocyanine analog IC3-dCTP and the rhodamine analog R6G-dCTP. It should be noted that these studies used normal deoxyribose sugars, and thus the ability of these analogs to be incorporated as chain terminators was not addressed.

Although ddNTPs are the dominant chain terminators utilized, other analogs have also been explored as chain terminators. The use of acyclo-nucleoside triphosphates (acyclo-NTPs) in chain terminator DNA sequencing by the Klenow fragment of DNA polymerase I and by AMV reverse transcriptase has been discussed (US Patent No. 5,558,991). Such acyclo derivatives substitute a 2-hydroxyethoxymethyl group for the 2'-deoxyribofuanosyl sugar normally present in dNTPs. Sequencing patterns produced by these two enzymes were found to be virtually identical for use of ddNTPs and acyclo-NTPs. However, approximately ten-fold higher concentrations of the acyclo derivatives were required to produce equivalent

patterns, indicating a greater discrimination against those compounds by the two enzymes tested. Thus, at least for these two enzymes, ddNTPs are favored substrates over acyclo-NTPs.

5 Incorporation of acyclo-NTPs has also been analyzed with Family B DNA polymerases, especially in light of the antiviral activity of selected acyclo derivatives, specially acyclovir. The mode of drug action, in part, is thought to be the preferential incorporation of the chain terminator acyclovir (9-(2-hydroxyethoxymethyl)guanine] triphosphate) by the viral as opposed to the
10 cellular DNA polymerase, human DNA polymerase alpha (Elion, *J. Antimicrob. Chemother.* **12** Suppl. B:9-17 (1983)). This conclusion is tempered by the observation that inhibition by acycloguanosine has both competitive and non-competitive components. Evidence has been proffered that the Family B herpes virus type 2 (HSV-2) and human cytomegalovirus
15 (HCMV) DNA polymerases have a preference for insertion of acyclo-GTP over ddGTP (Reid, *et al.*, *J. Biol. Chem.* **263**:3898-3904 (1988)). The same report also indicates a strong preference by human DNA polymerase alpha (also a Family B DNA polymerase) for insertion of ddGTP over acyclo-GTP. The contrasting behaviors of these three Family B DNA polymerases,
20 combined with the complex inhibition patterns observed for the viral polymerases, makes *a priori* predictions difficult for other Family B DNA polymerases as far as the relative incorporation of ddNTPs and acyclo-NTPs are concerned.

25 SUMMARY OF THE PRESENT INVENTION

 The present invention is directed toward improving the efficiency of chain terminator incorporation by Family B archaeon DNA polymerases. Previously, the low efficiency of ddNTP, and more especially dye-labeled
30 ddNTP, incorporation has limited the usefulness of this group of DNA polymerases in protocols requiring chain terminator incorporation.

 Several innovations are exploited in novel combinations in the present invention to overcome previously-noted limitations in chain terminator
35 incorporation. In accordance with the present invention, derivatized ddNTP

terminators are identified that are more efficiently incorporated than the corresponding underivatized ddNTPs. Methods are delineated to identify additional compounds of this type. Such compounds offer a marked advantage over previously tested dye-labeled ddNTPs whose incorporation was disfavored.

In certain preferred embodiments, acyclo-NTP terminators are found to be more efficiently incorporated than the corresponding ddNTPs. As with ddNTPs, incorporation of these acyclo-NTPs can be enhanced by specific base adducts.

In other preferred embodiments, incorporation of acyclo-NTPs and of derivatized ddNTPs and acyclo-NTPs is further enhanced by use of DNA polymerase variants.

Each embodiment confers a significant advantage in terminator incorporation, most strongly seen when the various embodiments are combined in a single reaction. In a most preferred embodiment, a variant DNA polymerase is used to incorporate a derivatized acyclo-NTP, using polymerase variants and derivatized terminators typified in the present invention. This novel arrangement provides a vast increase in terminator incorporation over that previously reported.

The efficient production of chain terminator products has obvious application in DNA sequence determination. This arises not only in traditional chain terminator sequencing, but also in automated procedures where detection is *via* incorporation of dye-labeled terminators. The present invention is applicable to both long range DNA sequence determination where hundreds of base pairs of contiguous sequence are revealed, and to short range sequencing, defining as little as one base pair of sequence. In the case of short range sequencing, the present invention is useful in analyzing sequence polymorphisms, for example in genetic testing and screening for specific single nucleotide polymorphisms (SNPs). Characterization of SNPs can be either by virtue of molecular weight or label incorporation, in either case accommodated by methods described in the present invention.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows the incorporation of modified ddCTP bases by Vent® (exo-) and Thermo Sequenase™ DNA polymerases. Extension of a [³²P]-labeled primer on an M13mp18 single-stranded substrate was examined in the presence of a 1:1 ratio or 1:10 ratio of analog to dNTP. A reaction containing a 1:1 ratio of unmodified ddCTP to dNTP is used for reference in the first lane. Lanes marked "dNTP" are control reactions performed in the absence of terminators. A, Vent® (exo-). B, Thermo Sequenase™.

Figure 2 compares dye-labeled ddCTP and dye-labeled acyclo-CTP incorporation by Vent® (exo-) and Thermo Sequenase™ DNA polymerases. Extension of a [³²P]-labeled primer on an M13mp18 single-stranded substrate was examined in the presence of a 1:1 ratio or 1:10 ratio of analog to dNTP. In each panel, a reaction containing a 1:1 ratio of unmodified ddCTP to dNTP is used for reference in the first lane. A, Vent® (exo-). B, Thermo Sequenase™.

Figure 3 demonstrates the incorporation efficiency of ROX-acyclo-CTP by Vent® (exo-), Deep Vent® (exo-), Pfu (exo-) and 9°N™ (exo-) DNA polymerases. Numbers refer to the ratio of ROX-acyclo-CTP : dCTP in the reaction mixture. The lane labeled "dNTP" illustrates a control reaction not containing terminators.

Figure 4 shows the incorporation of modified ddCTP bases by Vent® (exo-)/A488L DNA polymerase. Extension of a [³²P]-labeled primer on an M13mp18 single-stranded substrate was examined in the presence of a 1:1 ratio or 1:10 ratio of analog to dNTP. In each panel, a reaction containing a 1:1 ratio of unmodified ddCTP to dNTP is used for reference in the first lane and a reaction containing dNTP but lacking terminators is also shown.

Figure 5 compares the incorporation efficiency of ROX-ddCTP by Vent® (exo-), Vent® (exo-)/A488L and Vent® (exo-)/Y499L DNA polymerases. Numbers refer to the ratio of ROX-ddCTP : dCTP in the

reaction mixture. The reaction in the lane labeled "dNTP" contains no chain terminators.

Figure 6 compares the incorporation efficiency of ROX-ddCTP and ddCTP by Vent® (exo-), Vent® (exo-)/A488L, 9°N™ (exo-) and 9°N™ (exo-)/A485L DNA polymerases. Numbers refer to the ratio of ROX-ddCTP:dCTP or ddCTP:dCTP in the reaction mixture.

Figure 7 compares incorporation of ROX, IRD700 and TAMRA dye-labeled ddCTP and acyclo-CTP by Vent® (exo-)/A488L DNA polymerase. Numbers refer to the ratio of ROX-ddCTP:dCTP in the reaction mixture.

Figure 8 compares incorporation of ddCTP and IRD700, ROX and TAMRA dye-labeled acyclo-CTP by Vent® (exo-)/A488L, 9°N™ (exo-)/A485L and Thermo Sequenase™ DNA polymerases. The terminator was present in a 1:1 ratio with dCTP in all cases. Lanes marked dNTP delineate reactions without added terminators.

Figure 9 compares incorporation of ddGTP and acyclo-GTP by Thermo Sequenase™ and 9°N™ (exo-)/A485L DNA polymerases. Numbers refer to the ratio of terminator : dGTP in the reaction mixture.

Figure 10 illustrates the output of an ABI377 automated DNA sequencer with samples generated with either 9°N™ (exo-)/A485L DNA Polymerase or AmpliTaq® DNA Polymerase, FS. The DNA sequence along the top line is the consensus sequence from the two unedited traces, while those above the traces are sequences assigned by AutoAssembler software (Perkin-Elmer Corp.).

DETAILED DESCRIPTION OF THE PRESENT INVENTION

In accordance with the present invention, three innovations are utilized to facilitate incorporation of modified nucleotides by DNA polymerases. Each of these three elements makes a significant individual contribution to the

-10-

efficiency of incorporation, and can act in concert with the other elements to further enhance incorporation of chain terminators. These elements are, (1) functionalities whose attachment to nucleotide bases can enhance incorporation of that base relative to the naturally-occurring base, and methods to identify such compounds, (2) acyclo-NTPs, based on the discovery that such compounds are more readily incorporated than corresponding ddNTP derivatives by archaeon DNA polymerases and (3) identification and use of archaeon DNA polymerases and polymerase variants with enhanced ability to incorporate nucleotides with modified sugars, specifically chain terminators such as ddNTPs and acyclo-NTPs.

Identifying DNA polymerases with similar properties

As mentioned above, DNA polymerases can be categorized into three families, with enzymes such as Vent® falling into Family B. DNA polymerases within a family can be further subdivided into groups with similar features. Such groupings can be made by several criteria. First, through analytical methods that detect the degree of homology in the underlying nucleic acid sequences encoding the gene. Such similarities are sufficient in many cases to isolate similar genes from alternate organisms, and has been used to discover new archaeon Family B DNA polymerases, as described in US Patent No. 5,500,363. In that invention, specific DNA probes and hybridization conditions are described to allow for detection by Southern Blot, and isolation of such similar DNA polymerases. The DNA fragment encoding the DNA polymerase was identified as that hybridizes in a Southern blot to an isolated DNA fragment selected from the group consisting of a DNA fragment having nucleotides 1-1274 of SEQ ID NO:4, a DNA fragment having nucleotides 291-1772 of SEQ ID NO:4, a DNA fragment having nucleotides 3387-3533 of SEQ ID NO:4, a DNA fragment having nucleotides 4704-5396 of SEQ ID NO:4, and a DNA fragment having nucleotides 4718-5437 of SEQ ID NO:4, wherein hybridization is conducted under the following conditions: a) hybridization: 0.75 M NaCl, 0.15 M Tris, 10 mM EDTA, 0.1% sodium pyrophosphate, 0.1% sodium lauryl sulfate, 0.03% BSA, 0.03% Ficoll 400, 0.03% PVP and 100 µg/ml boiled calf thymus DNA at 50°C for about 12 hours and; b) wash: 3X30 minutes with

-11-

0.1X SET, 0.1% SDS, 0.1% sodium pyrophosphate and 0.1 M phosphate buffer at 45°C.

SEQ ID NO:4

5

GAATTCGCGATAAAATCTATTTTCTTCCTCCATTTTTCATTTTCAAAAACGTAAGCATGA 60
GCCAAACCTCTCGCCCTTTCTCTGTCTTCCCGCTAACCTCTTGAAAACCTCTCTCCAAA 120
GCATTTTTTGATGAAAGCTCACGCTCCTCTATGAGGGTCAGTATATCTGCAATGAGTTCG 180
TGAAGGGTTATTCTGTAGAACAACTCCATGATTTTTCGATTTGGATGGGGGTTTAAAAATT 240
TGGCGGAACTTTATTTAATTTGAACCTCAGTTTATATCTGGTGGTATTTATGATACTGG 300
ACACTGATTACATAACAAAAGATGGCAAGCCTATAATCCGAATTTTAAAGAAAGAGAACG 360
GGGAGTTTAAAATAGAACTTGACCCCTCATTTTCAGCCCTATATATATGCTCTTCTCAAAG 420
ATGACTCCGCTATTGAGGAGATAAAGGCAATAAAGGGCGAGAGACATGGAAAACTGTGA 480
GAGTGCTCGATGCAGTGAAAGTCAGGAAAAAATTTTGGGAAGGGAAGTTGAAGTCTGGA 540
AGCTCATTTTCGAGCATCCCCAAGACGTTCCAGCTATGCGGGGCAAAATAAGGGAACATC 600
CAGCTGTGGTTGACATTTACGAATATGACATACCCTTTGCCAAGCGTTATCTCATAGACA 660
AGGGCTTGATTCCCATGGAGGGAGACGAGGAGCTTAAGCTCCTTGCCTTTGATATTGAAA 720
CGTTTTATCATGAGGGAGATGAATTTGGAAAGGGCGAGATAATAATGATTAGTTATGCCG 780
ATGAAGAAGAGGCCAGAGTAATCACATGGAAAAATATCGATTTGCCGTATGTCGATGTTG 840
TGTCCAATGAAAGAGAAATGATAAAGCGTTTGTTCAGTTGTTAAAGAAAAAGACCCCG 900
ATGTGATAATAACTTACAATGGGGACAATTTTGATTTGCCGTATCTCATAAAACGGGCAG 960
AAAAGCTGGGAGTTTCGGCTTGTCTTAGGAAGGGACAAAGAACATCCCGAACCCAAGATTC 1020
AGAGGATGGGTGATAGTTTTGCTGTGGAAATCAAGGGTAGAATCCACTTTGATCTTTTCC 1080
CAGTTGTGCGAAGGACGATAAACCTCCCAACGTATACGCTTGAGGCAGTTTATGAAGCAG 1140
TTT TAGGAAAAACCAAAGCAAATTAGGAGCAGAGGAAATTGCCGCTATATGGGAAACAG 1200
AAGAAAGCATGAAAAAACTAGCCCAGTACTCAATGGAAGATGCTAGGGCAACGTATGAGC 1260
TCGGGAAGGAATCTTCCCATGGAAGCTGAGCTGGCAAAGCTGATAGGTCAAAGTGTAT 1320
GGGACGTCTCGAGATCAAGCACCGGCAACCTCGTGGAGTGGTATCTTTTAAGGGTGGCAT 1380
ACGCGAGGAATGAACCTGCACCGAACAAACCTGATGAGGAAGAGTATAAACGGCGCTTAA 1440
GAACAACCTTACCTGGGAGGATATGTAAAAGAGCCAGAAAAAGGTTTGTGGGAAAATATCA 1500
TTTATTTGGATTTCCGCAGTCTGTACCCTTCAATAATAGTTACTCACAACGTATCCCCAG 1560
ATACCCTTGAAAAAGAGGGCTGTAAGAATTACGATGTTGCTCCGATAGTAGGATATAGGT 1620
TCTGCAAGGACTTTCCGGGCTTTATTCCCTCCATACTCGGGGACTTAATTGCAATGAGGC 1680
AAGATATAAAGAAGAAAATGAAATCCACAATTGACCCGATCGAAAAGAAAATGCTCGATT 1740
ATAGGCAAAGGGCTATTAAATTGCTTGCAAACAGCATCTTACCCAACGAGTGGTTACCAA 1800
TAATTGAAAATGGAGAAATAAAATTCGTGAAAATTGGCGAGTTTATAAACTCTTACATGG 1860
AAAAACAGAAGGAAAACGTTAAACAGTAGAGAATACTGAAGTTCTCGAAGTAAACAACC 1920

35

-12-

TTTTTGCATTCTCATTCAACAAAAAATCAAAGAAAGTGAAGTCAAAAAAGTCAAAGCCC 1980
TCATAAGACATAAGTATAAAGGGAAAGCTTATGAGATTGAGCTTAGCTCTGGTAGAAAAA 2040
TTAACATAACTGCTGGCCATAGTCTGTTTACAGTTAGAAATGGAGAAATAAAGGAAGTTT 2100
CTGGAGATGGGATAAAAAGAAGGTGACCTTATTGTAGCACCAAAGAAAATTAAACTCAATG 2160
5 AAAAAGGGGTAAGCATAAACATTCCCGAGTTAATCTCAGATCTTTCCGAGGAAGAAACAG 2220
CCGACATTGTGATGACGATTTGAGCCAAGGGCAGAAAGAACTTCTTTAAAGGAATGCTGA 2280
GAACTTTAAGGTGGATGTTTGGAGAAGAAAATAGAAGGATAAGAACATTTAATCGCTATT 2340
TGTTCCATCTCGAAAACTAGGCCTTATCAAACACTGCCCCGCGGATATGAAGTTACTG 2400
ACTGGGAGAGATTAAAGAAATATAAACAACTTTACGAGAAGCTTGCTGGAAGCGTTAAGT 2460
10 ACAACGGAAACAAGAGAGAGTATTTAGTAATGTTCAACGAGATCAAGGATTTTATATCTT 2520
ACTTCCACAAAAAGAGCTCGAAGAATGGAAAATTGGAACCTCTCAATGGCTTTAGAACGA 2580
ATTGTATTCTCAAAGTCGATGAGGATTTTGGGAAGCTCCTAGGTACTATGTTAGTGAGG 2640
GCTATGCAGGTGCACAAAAAATAAACTGGTGGTATCAGTTATTCGGTGAAGCTTTACA 2700
ATGAGGACCTAATGTTCTTGAGAGCATGAAAAATGTTGCAGAAAAATTCTTTGGCAAGG 2760
15 TTAGAGTTGACAGAAATTGCGTAAGTATATCAAAGAAGATGGCATACTTAGTTATGAAAT 2820
GCCTCTGTGGAGCATTAGCCGAAAACAAGAGAATTCCTTCTGTTATACTCACCTCTCCCG 2880
AACCGGTACGGTGGTCATTTTGTAGAGGCGTATTTTACAGGCGATGGAGATATACATCCAT 2940
CAAAAAGGTTTAGGCTCTCAACAAAAAGCGAGCTCCTTGCAAATCAGCTTGTGTTCTTGC 3000
TGAACCTTTTGGGAATATCCTCTGTAAAGATAGGCTTTGACAGTGGGGTCTATAGAGTGT 3060
20 ATATAAATGAAGACCTGCAATTTCCACAAACGTCTAGGGAGAAAAACACATACTACTCTA 3120
ACTTAATTTCCCAAAGAGATCCTTAGGGACGTGTTTGGAAAAGAGTTCCAAAAGAACATGA 3180
CGTTCAAGAAATTTAAAGAGCTTGTGACTCTGGAAAACCTAACAGGGAGAAAGCCAAGC 3240
TCTTGGAGTTCTTCATTAATGGAGATATTGTCTTGACAGAGTCAAAAAGTGTAAAGAAA 3300
AGGACTATGAAGGGTATGTCCTATGACCTAAGCGTTGAGGATAACGAGAACTTTCTTGTG 3360
25 GTTTTGGTTTGTCTATGCTCACAACAGCTATTACGGCTATATGGGGTATCCTAAGGCAA 3420
GATGGTACTCGAAGGAATGTGCTGAAAGCGTTACCGCATGGGGGAGACACTACATAGAGA 3480
TGACGATAAGAGAAATAGAGGAAAAGTTCGGCTTTAAGGTTCTTTATGCGGACAGTGTCT 3540
CAGGAGAAAGTGAGATCATAATAAGGCAAAACGGAAAGATTAGATTTGTGAAAATAAAGG 3600
ATCTTTTCTCTAAGGTGGACTACAGCATTGGCGAAAAAGAATACTGCATTCTCGAAGGTG 3660
30 TTGAAGCACTAACTCTGGACGATGACGGAAAGCTTGTCTGGAAGCCCGTCCCTACGTGA 3720
TGAGGCACAGAGCGAATAAAAGAATGTTCCGCATCTGGCTGACCAACAGCTGGTATATAG 3780
ATGTTACTGAGGATCATTCCTCATAGGCTATCTAAACACGTCAAAAACGAAAAC TGCCA 3840
AAAAATCGGGGAAAGACTAAAGGAAGTAAAGCCTTTTGAATTAGGCAAAGCAGTAAAT 3900
CGCTCATATGCCCAAATGCACCGTTAAAGGATGAGAATACCAAAACTAGCGAAATAGCAG 3960
35 TAAAATTCTGGGAGCTCGTAGGATTGATTGTAGGAGATGGAACTGGGGTGGAGATTCTC 4020
GTTGGGCAGAGTATTATCTTGGACTTTCAACAGGCAAAGATGCAGAAGAGATAAAGCAA 4080
AACTTCTGGAACCCCTAAAACTTATGGAGTAATCTCAAACATTATACCAAAAAACGAGA 4140
AAGGGGACTTCAACATCTTGGCAAAGAGCCTTGTAAGTTTATGAAAAGGCACTTTAAGG 4200

-13-

5 ACGAAAAAGGAAGACGAAAAATTCCAGAGTTCATGTATGAGCTTCCGGTTACTTACATAG 4260
 AGGCATTTCTACGAGGACTGTTTTTCAGCTGATGGTACTGTAACATCAGGAAGGGAGTTC 4320
 CAGAGATCAGGCTAACAAACATTGATGCTGACTTTCTAAGGGAAGTAAGGAAGCTTCTGT 4380
 GGATTGTTGGAATTTCAAATTCAATATTTGCTGAGACTACTCCAAATCGCTACAATGGTG 4440
 10 TTTCTACTGGAACCTACTCAAAGCATCTAAGGATCAAAAATAAGTGGCGTTTTGCTGAAA 4500
 GGATAGGCTTTTTAATCGAGAGAAAGCAGAAGAGACTTTTAGAACATTTAAAATCAGCGA 4560
 GGGTAAAAAGGAATACCATAGATTTTGGCTTTGATCTTGTGCATGTGAAAAAAGTCGAAG 4620
 AGATACCATACGAGGGTTACGTTTATGACATTGAAGTCGAAGAGACGCATAGGTTCTTTG 4680
 CAAACAACATCCTGGTACACAATACTGACGGCTTTTATGCCACAATACCCGGGGAAAAGC 4740
 15 CTGAACCTATTAAAAAGAAAGCCAAGGAATTCCTAAACTACATAAACTCCAAACTTCCAG 4800
 GTCTGCTTGAGCTTGAGTATGAGGGCTTTTACTTGAGAGGATTCTTTGTTACAAAAAAGC 4860
 GCTATGCAGTCATAGATGAAGAGGGCAGGATAACAACAAGGGGCTTGGAAGTAGTAAGGA 4920
 GAGATTGGAGTGAGATAGCTAAGGAGACTCAGGCAAAGGTTTTAGAGGCTATACTTAAAG 4980
 AGGGAAGTGTTGAAAAAGCTGTAGAAGTTGTTAGAGATGTTGTAGAGAAAAATAGCAAAAT 5040
 20 ACAGGGTTCCACTTGAAAAGCTTGTTATCCATGAGCAGATTACCAGGGATTTAAAGGACT 5100
 ACAAAGCCATTGGCCCTCATGTCGCGATAGCAAAAAGACTTGCCGCAAGAGGGGATAAAAG 5160
 TGAAACCGGGCACAATAATAAGCTATATCGTTCTCAAAGGGAGCGGAAAGATAAGCGATA 5220
 GGGTAATTTTACTTACAGAATACGATCCTAGAAAACACAAGTACGATCCGGACTACTACA 5280
 TAGAAAACCAAGTTTTGCCGGCAGTACTTAGGATACTCGAAGCGTTTGGATACAGAAAGG 5340
 25 AGGATTTAAGGTATCAAAGCTCAAACAAACCGGCTTAGATGCATGGCTCAAGAGGTAGC 5400
 TCTGTTGCTTTTTAGTCCAAGTTTCTCCGCGAGTCTCTCTATCTCTCTTTTGTATTCTGC 5460
 TATGTGGTTTTTCATTCATTAAGTAGTCCGCCAAAGCCATAACGCTTCCAATTCCAAA 5520
 CTTGAGCTCTTTCCAGTCTCTGGCCTCAAATTCATCCATGTTTTTGGATCGTCGCTTCT 5580
 CCCTCTTCTGCTAAGCCTCTCGAATCTTTTTCTTGCGAAGAGGTGTACAGCTATGATGAT 5640
 30 TATCTCTTCTCTGGAAACGCATCTTTAAACGTCTGAATTTTCATCTAGAGACCTCACTCC 5700
 GTCGATTATAACTGCCTTGTAATTCTTTAGTAGTTCTTTTACCTTTGGGATCGTTAATTT 5760
 TGCCACGGCATGTGCCCAAGCTCCTGCCTAAGCTGAATGCTCACACTGTTTCATACCTTC 5820
 GGGAGTTCTTGGGATCC 5837

30 In a similar vein, analytical methods can also be used to discover and identify proteins with similar amino acid sequences, for example by using antibodies raised to a first DNA polymerase to identify other related proteins based on cross-reactivity (US Patent No. 5,500,363).

35 A second method of grouping is by the degree of identity and/or similarity between the primary amino acid sequence of the polymerases, which the worker skilled in the art will recognize as also being correlated to the

underlying gene coding sequence. This method of analysis relies on sequence alignments rather than physical characterization. Several computer programs have been devised to make this comparison between proteins, one of which is BLAST (Altschul, *et al.* *Nucleic Acids Res.* **25**:3389-3402 (1997); Tatusova, *et al.*, *FEMS Microbiol Lett.* **174**:247-250 (1999)). Alignments obtained with such programs typically report the percentage of sequence identity and of sequence similarity between two test sequences, although other statistical measures of identity and similarity are also available and can be relied upon. In addition to this global measure of sequence similarity, proteins can also display sequence similarity over short stretches of primary amino acid sequence. While not wishing to be bound by theory, these patches of similarity are thought to occur most often at essential protein interfaces, such as those involved in catalysis, substrate binding or protein-protein recognition. As such, the degree of sequence similarity, particularly in conserved sequence motifs, is predictive of the degree to which the proteins will behave similarly in both physical properties and catalytic function. For example, mutation of the motif associated with 3'-5' exonuclease activity of DNA polymerases (Bernad, *et al.*, *Cell* **59**:219-228 (1989)) has been shown to abolish this activity in a variety of polymerases (Derbyshire, *et al.*, *Science* **240**:199-201 (1988)). Example 3 illustrates BLAST-derived sequence identity information for selected archaeon DNA polymerases.

Finally, groupings can be defined by functional similarity, assessed by biochemical assays of such features as kinetic parameters (*e.g.*, K_m and turnover number), propensity to insert modified nucleotides, template specificity, and sensitivity to changes in reaction conditions such as pH, temperature, salt types and composition, and cofactors (*e.g.*, Mg^{2+}). For the purposes of this invention, the most important grouping is by functional assays of the polymerases, *i.e.* the ability to efficiently incorporate the modified bases described herein.

As the examples will show, the above groupings are inter-related. Thus, DNA polymerases grouped together by sequence similarities, both nucleic acid and amino acid, also tend to have similar biochemical characteristics. Thus, a reasonable prediction is that DNA polymerases

-15-

showing a greater degree of similarity to those archaeon DNA polymerases in the examples will be most likely to function in the invention described herein.

5 ***DNA polymerase variants with diminished exonuclease activity***

Two general classes of archaeon DNA polymerase variants are utilized in the present invention. First, exonuclease-deficient (exo-) variants. A number of DNA polymerases possess a 3-5' exonuclease activity, including
10 the Family B DNA polymerases identified in archaea. One function of this activity is "proofreading," wherein the polymerase can remove 3' nucleotides before proceeding with polymerization. Incorrectly base-paired, or aberrant nucleotides are preferentially removed by this activity, increasing the fidelity of replication (Kornberg, *DNA Replication*, W.H. Freeman and Company,
15 San Francisco, p. 127 (1980)). While not wishing to be bound by theory, modified nucleotides might reasonably be expected to sensed as aberrant, and, even if incorporated, be subject to removal by this activity. To avoid this possibility, variants have been created that lack or have diminished exonuclease activity (Vent® DNA polymerase: Kong, *et al.*, *supra* (1993);
20 US Patent No. 5,352,778; *Pyrococcus furiosus* (*Pfu*) DNA polymerase: US Patent No. 5,489,523; *Tba* DNA polymerase: US Patent No. 5,882,904; Deep Vent® DNA polymerase: US Patent No. 5,834,285, MA; 9°N™ DNA polymerase: Southworth, *et al.*, *Proc. Natl. Acad. Sci. USA* **93**:5281-5285 (1996); KOD DNA polymerase: US Patent No. 6,008,025). In each of
25 these cases exonuclease activity was diminished by creating polymerases with specific variations within a common, recognized amino acid sequence motif, enabling the skilled artisan to predict where similar changes could be made in other DNA polymerases to similarly modulate exonuclease activity. One skilled in the art will appreciate the possibility that the exonuclease-
30 deficient forms may not be absolutely required in this application, as suggested in US Patent No. 5,945,312 for the Family B DNA polymerase derived from bacteriophage T4. The second general class of DNA polymerases variants will be described below.

Incorporation of dye-labeled nucleotides by archaeon DNA polymerases

At present, it is difficult to predict on structural or chemical grounds which dye-substituted molecules will be readily incorporated by archaeon DNA polymerases. A limited literature has reported incorporation of nucleotides with dye-labeled bases by archaeon DNA polymerases. US Patent No. 5,723,298 claims incorporation of IRD40 dATP by Vent® DNA polymerase, although measures of the efficiency of incorporation were not presented. Recently, Ilsey and Buzby (Ilsey and Buzby, *supra.*) identified dye-labeled dCTP analogs whose incorporation was enhanced relative to underivatized dCTP by Vent® (exo-) DNA polymerase, as evident in comparative incorporation assays. Ilsey and Buzby noted in the series of compounds tested that the most hydrophobic cyanine dye was the preferred substrate for polymerization (Ilsey and Buzby, *supra.*). However, due to the limited number of dyes tested, there is little predictive power in this observation.

Incorporation of dye-substituted chain terminators, however, was not examined in the above studies, allowing uncertainty as to whether these results can be extended to instances where the dye is linked to a chain terminator. In fact, initial studies of Vent® (exo-) DNA polymerase suggested that dye-terminator incorporation by Vent® DNA polymerase was not efficient ("CircumVent®: Questions and Answers," *The NEB Transcript*, September 1992, p. 12-13). Additionally, US Patent No. 5,882,904 reports Deep Vent® (exo-) and *Tba* (exo-) DNA polymerases incorporate FL-ddNTPs many times less efficient than a modified form of Taq DNA polymerase (KlenTaq). Both of these reports leave significant questions as to whether dye-terminators themselves are disfavored, and if alternative dyes or terminator functionalities can influence the observed discrimination against incorporation.

Dye-terminators that are efficiently incorporated by archaeon DNA polymerases

In order to determine the extent of dye-terminator incorporation by archaeon DNA polymerases, the titration assay described by Gardner and Jack (*supra.*) was used (Example 1). In this assay, the efficiency of incorporation of chain terminator nucleotides is judged by the size distribution of reaction products in a polymerization reaction. As the efficiency of chain-terminator incorporation increases, the average reaction product size decreases because polymerization is more often halted by terminator addition. By comparing the amount of terminator required to give the same spectrum of reaction products, the relative efficiency of incorporation of the test compounds with the different polymerases can be determined. Initial determinations compared the ability of Thermo Sequenase™ (Amersham Pharmacia Biotech, Piscataway, NJ) and Vent® (exo-) (New England Biolabs, Beverly, MA) to incorporate a variety of dye-substituted ddCTP analogs, comparing their incorporation to underivatized ddCTP. Dye terminators were obtained from NEN Life Sciences (Boston, MA), either as commercial products or as evaluation samples (Table 2).

Three classes of dye terminators could be identified with Vent® (exo-) DNA polymerase based on the patterns of termination products produced (Example 2). In the first class banding patterns for ddCTP and the analog had similar concentration dependencies, indicating that the modified base was incorporated no better than the corresponding ddCTP (*e.g.*, IRD40 ddCTP; Figure 1A). In the second class incorporation of the dye-substituted base was less than that observed with the normal ddCTP as indicated by a dominance of higher molecular weight bands at fixed terminator concentration (*e.g.*, JOE ddCTP; Figure 1A). In the final class of analogs the distribution of terminated products was shifted to lower molecular weights, indicating an increased incorporation relative to the corresponding ddCTP substrate (*e.g.*, ROX ddCTP, TAMRA ddCTP, BODIPY® TR ddCTP and BODIPY® TMR ddCTP ; Figure 1A). In this last set of analogs, the presence of the dye enhanced incorporation of the terminator base relative to the parent base ddCTP. Importantly, when appropriate concentrations of analog were

compared, a similar band pattern emerged, indicating no loss of base specificity in the insertion of the analogs. Although a limited number of dyes were analyzed by this method, the efficiency of incorporation of alternate dyes could easily be evaluated using the same assay system in conjunction with the desired DNA polymerase. Of course, alternate assays that compare the relative ability of modified nucleotides to be incorporated could also be employed.

Relative incorporation of dye-labeled ddNTP and acyclo-NTP derivatives

The above experiments identified dyes that, when coupled to ddNTPs, enhanced nucleotide incorporation. Acyclo-NTP derivatives have been used as terminators, although the literature does not report instances of their use with archaeon DNA polymerases. To test whether this alternate terminator might also function in this system, incorporation of dye-acyclo-CTP derivatives were tested using a titration assay (Example 5). In each case tested, acyclo derivatives were more efficiently incorporated by Vent® (exo-) DNA polymerase than those of the corresponding ddNTP (ROX-acyclo-CTP, IRD700-acyclo-CTP and TAMRA-acyclo-CTP; Figure 2A), indicating that the acyclo chain terminator was even more effectively incorporated than the dideoxy analog. The hierarchy of which dye terminators were more efficiently incorporated was identical for ddNTP and acyclo-NTP derivatives, serving as additional confirmation that the acyclo-NTPs are better incorporated than ddNTPs by the archaeon DNA polymerases (see also Example 12). In contrast, incorporation of acyclo derivatives by Thermo Sequenase™ did not exceed ddNTP incorporation levels, and in most cases was significantly lower (Figure 2B).

The similarity in amino acid sequence of Family B archaeon DNA polymerases suggests that these enzymes might share similar dye-terminator incorporation properties. Table 3 is a sampling of other Family B DNA polymerases from sequence databases, comparing the primary amino acid sequence similarity for the proteins. As described above, those with

greatest similarity to the tested enzymes are considered most likely to share the described incorporation properties.

Accordingly, terminator titrations assays were repeated using an expanded set of archaeon DNA polymerases, specifically Vent® (exo-) (New England Biolabs, Beverly, MA), Deep Vent® (exo-) (New England Biolabs, Beverly, MA), Pfu (exo-) (Stratagene, La Jolla, CA) and 9N™ (exo-) (Example 4). The pattern of incorporation preference was identical for these four enzymes (see Example 6, Figure 3), each demonstrating more efficient incorporation of dye-acyclo-NTPs than the corresponding dye-ddNTPs. Thus, the dye-terminator incorporation properties of one enzyme should be predictive of the incorporation properties of other members of this set.

The similarity of incorporation patterns with these selected enzymes suggests that not only these archaeon DNA polymerases, but a larger family of DNA polymerases could share the ability to incorporate acyclo to a greater extent than dideoxy terminators. Since *Pfu*, Deep Vent® and 9°N™ DNA polymerases have greater than about 70% sequence identity with Vent DNA polymerase, other enzymes with equivalent or greater identity can reasonably be expected to perform as Vent® (exo-) DNA polymerase in this invention. Notably, those enzymes for which no significant sequence similarity is found (*i.e.*, Family A DNA polymerases such as Taq) do not perform in similar ways in the current invention. This fact leads us to believe that archaeon enzymes showing intermediate identity, namely those between about 20 and 70% identity, are reasonable candidates for this invention. Indeed, the Family B herpes simplex virus type 2 and human cytomegalovirus DNA polymerases in that range of sequence identity have been reported to incorporate acyclo-GTP to a greater extent than ddGTP (Reid, *et al.*, *J. Biol. Chem.* **263**:3898-3904 (1988)). Some caution must be employed in this regard as the same report indicates human DNA polymerase alpha (27% sequence identity) incorporates ddGTP to a much greater extent than acyclo-GTP (Reid, *et al.*, *supra*). While not wishing to be bound by theory, we anticipate the lack of utility of the human DNA polymerase alpha in this invention arises from the evolutionary distance

Table 3

Comparison Of Region III Of A Variety Of Family B DNA Polymerases

	<u>DNA polymerase source</u>	<u>Accession</u>	<u>Motif B</u>	<u>Similarity</u>	<u>SEQ ID NO:</u>
5	Vent® (Thermococcus litoralis)	AAA72101	AIKLLANSYYGYMGY	100/100/0	SEQ ID NO:5
	Deep Vent® (Pyrococcus sp. GB-D)	AAA67131	AIKILANSYYGYGY	77/88/0	SEQ ID NO:6
	9 ^N 7™ (Thermococcus sp.)	pdb 1QHT A	AIKILANSFYGYGY	77/87/0	SEQ ID NO:7
	Pyrococcus furiosus	S35543	AIKLLANSFYGYGY	74/87/0	SEQ ID NO:8
	Thermococcus fumicolans	CAA93738	AIKILANSFYGYGY	78/88/0	SEQ ID NO:9
10	Thermococcus gorgonarius	P56689	AIKILANSFYGYGY	79/89/0	SEQ ID NO:10
	Thermococcus sp. (TY)	CAA73475	AVKLLANSYYGYMGY	94/98/0	SEQ ID NO:11
	Pyrococcus abyssi	CAA90888	AIKILANSYYGYGY	75/86/0	SEQ ID NO:12
	Pyrococcus glycoaorans	CAB81809	AIKILANSYYGYGY	77/88/0	SEQ ID NO:13
	Pyrococcus horikoshii	BAA31074	AIKILANSYYGYGY	75/87/0	SEQ ID NO:14
15	Pyrococcus sp. (GE23)	Z54173	AIKILANSYYGYGY	76/87/0	SEQ ID NO:15
	Pyrococcus sp. (KOD1)	S71551	AIKILANSYYGYGY	78/87/0	SEQ ID NO:16
	Pyrococcus woesei	AAB67984	AIKLLANSFYGYGY	74/87/0	SEQ ID NO:17
	Archaeoglobus fulgidus	AAB90741	TLKVLTNFSFGYMGW	37/56/11	SEQ ID NO:18
	Cenarchaeum symbiosum	AAC62689	ALKVVLNASYGVMDGA	26/43/11	SEQ ID NO:19
20	Methanococcus jannaschi	AAB98889	SIKILANSVYGYLAF	46/65/4	SEQ ID NO:20

Table 3 (continued)

<u>DNA polymerase source</u>	<u>Accession</u>	<u>Motif B</u>	<u>Similarity</u>	<u>SEQ ID NO:</u>
Methanococcus voltae	AAA72443	SIKVLANSHYGYLAF	37/58/9	SEQ ID NO:21
Pyrodictium occultum	BAA07580	ALKVLANASYGYMGW	34/54/10	SEQ ID NO:22
Sulfurisphaera ohwakuensis	BAA23994	AMKVFINATYGVFGA	26/42/17	SEQ ID NO:23
Sulfolobus acidocaldarius	AAC44598	AMKVFINATYGVFGA	23/41/16	SEQ ID NO:24
Sulfolobus solfataricus	S23019	AMKVFINATYGVFGA	26/41/18	SEQ ID NO:25
Herpesvirus	DJBE21	AIKVV CNSVYGFTGV	22/41/16	SEQ ID NO:26
Human herpesvirus 2	NP_044550	AIKVV CNSVYGFTGV	22/41/16	SEQ ID NO:27
Human cytomegalovirus	P08546	ALKVTCNATYGFYGV	24/42/22	SEQ ID NO:28
human DNA polymerase alpha	NP_058633	ALKLTANSMYGCGLGF	27/46/8	SEQ ID NO:29
phage T4	AAD42468	NRKILINSLYGALGN	21/37/23	SEQ ID NO:30
Taq DNA polymerase	P19821	none		
Phage T7	P00581	none		

All listed protein sequences were accessed via the ncbi server at <http://www.ncbi.nlm.nih.gov> using the given accession numbers. Comparisons indicate % identity/% positives/% gaps, the output of the Blastp comparison of Vent® DNA polymerase with the listed DNA polymerase (using Blast 2; Tatusova, et al. *FEMS Microbiol Lett.* 174:247-250 (1999)) run on the site <http://www.ncbi.nlm.nih.gov/gorf/bl2.html> using the parameters: matrix 0BLOSUM62; gap open 11; gap extension 1; x_dropoff 50; expect 10; wordsize 3; filter off. Entries marked "none" returned "no significant similarity found" on Blastp processing. Motif B is as defined by Delarue, et al., *Protein Eng.* 3:461-467 (1990).

between humans and archaea, and thus should not discourage the worker skilled in the art from screening for the desired activity, for example by using the dye-terminator titrations described in this invention. We additionally note that the experiments by Reid, *et al.* did not encompass incorporation of dye-labeled substrates.

Vent DNA polymerase variants can increase dye-terminator incorporation

Archaeon DNA polymerase variants have been described that increase the incorporation efficiency for specific classes of chain terminators, namely ddNTPs and 3'-dNTPs. Variants of this type have been described in Vent® DNA polymerase (Gardner and Jack, *supra* (1999), *Pfu* DNA polymerase (US Patent No. 5,827,716) and *Tba* DNA polymerase (US Patent No. 5,882,904). In each of these examples, changes within a limited region of the protein primary sequence were made (Table 1), more specifically in Motif B (Delarue, *et al.*, *Protein Eng.* 3:461-467 (1990)). Most of these changes resulted in a modest (less than 5-fold) increase in incorporation of ddNTPs. Alterations corresponding to A488L in Vent® DNA polymerase produced the greatest increase (approximately 12-fold; Gardner and Jack, *supra.*), with lesser effects seen with a Y499L variant. It may be noted that the amino acid residues mutated in the above studies are absolutely conserved among the four polymerases tested, as well as among other archaeon DNA polymerases found in data bases (Table 3). It thus seems likely that similar mutations in related DNA polymerases will have similar effects.

Table 1

Primary sequence alignment for several archaeon DNA polymerases

<u>Vent®</u>	<u>9°N™</u>	<u>Tba</u>	<u>Pfu</u>
<u>A488</u>	<u>A485</u>	A485	A486
I489	I486	I486	I487
K490	K487	K487	K488
L491	I488	I488	L489
L492	L489	<u>L489</u>	L490
A493	A490	A490	<u>A491</u>
<u>Y499</u>	Y496	Y496	Y497

Rows present equivalent amino acid positions in the four DNA polymerases: *Thermococcus litoralis* (Vent®), *Thermococcus* sp. 9°N-7 (9°N™) and *Thermococcus barossii* (*Tba*). US Patent No. 5,827,716 discloses modification of residue A491 of *Pfu* DNA polymerase. US Patent No. 5,882,904 discloses modification of residue L489 of *Tba* DNA polymerase.

However, these studies did not address the effects of the cited mutations on dye-terminator incorporation, nor did they address incorporation of alternate terminators such as acyclo-NTPs. There was no teaching to suggest that the increased incorporation of ddNTPs by these variants could overcome the observed difficulty of incorporating dye-labeled ddNTPs ("CircumVent®: Questions and Answers," *supra*, US Patent No. 5,882,904), nor was there a teaching that dye-terminators could be incorporated by archaeon DNA polymerases more efficiently than the corresponding ddNTP. This invention goes beyond those previous studies in identifying dye-terminators better incorporated by archaeon Family B DNA polymerases, and additionally in using DNA polymerase variants to enhance incorporation.

The effects of the A488L mutation in dye-terminator incorporation was tested using the titration assay, comparing Vent® (exo-)/A488L and Vent® (exo-) DNA polymerases (Example 7; compare Figure 4 to Figures 1 and 3). Terminator incorporation was increased in the (exo-)/A488L variant when compared to the (exo-) polymerase. This increase was observed irrespective of whether the terminator was ddNTP or acyclo-NTP, and was observed with all dye-labels analyzed. When considered separately, the preference for acyclo-NTPs as opposed to ddNTPs was preserved, as was the relative preference for specific dyes noted above. Consequently, the increased incorporation reflects effects above and beyond those reported in the previous examples.

To broaden the applicability of this observation, the analogous 9°N™ (exo-) DNA polymerase variant A485L (Table 1) was created (Example 9) and employed in the same assays (see Examples 11 and 12). Relative incorporation of dye-terminators was similar to those observed with the

-24-

Vent® (exo-)/A488L DNA polymerase both with regards to terminator type (*i.e.*, ddNTP or acyclo-NTP; Figures 6, 8, and 9) and dye label (Figures 6 and 8), establishing the common effects of mutants at this locus among archaeon DNA polymerases.

Additional experiments with a second Vent® DNA polymerase variant, (exo-)/Y499L, displayed increased incorporation similar in nature to the (exo-)/A488L variant, demonstrating that increased incorporation was not limited to the (exo-)/A488L variant (Example 8; Figure 5). The worker skilled in the art will appreciate that variants analogous to those noted above in other archaeon DNA polymerases would be predicted to similarly enhance incorporation of comparable modified nucleotides. Additionally, other variants that facilitate incorporation of nucleotides with modified sugars might also be expected to operate in this invention, including but not necessarily limited to variants in residues corresponding to Vent® DNA polymerase residues A488 to Y499. For example, a variant corresponding to Vent® L492 has been described that increased ddNTP incorporation (US Patent No. 5,882,904). Titration assays, as described herein, could be used to verify the action of this and other similar variants in terminator incorporation.

Relative incorporation of ddGTP and acyclo-GTP

The experiments noted above gave very strong indications that the basis for enhanced dye-acyclo over dye-dideoxy terminator incorporation was primarily due to the modified sugar, and not the influence of the base. As a more direct test of this proposition, acyclo-GTP and ddGTP incorporation was compared for 9°N™ (exo-)/A485L and Thermo Sequenase™ DNA polymerases (Example 12). These two enzymes gave different responses, with Thermo Sequenase™ showing a preference for the ddGTP substrate, and 9°N™ (exo-)/A485L DNA polymerase showing a preference for acyclo-GTP. Results with the Family A polymerase Thermo Sequenase™ echo those reported in US Patent No. 5,558,991 for another Family A DNA polymerase, Klenow fragment, in the later case requiring a ten-fold higher acyclo-NTP concentration to see sequencing patterns identical to those obtained with ddNTP triphosphates. Thus, acyclo-NTPs are more efficient

terminators than ddNTPs with 9°N™ (exo-)/A485L, and by extension Family B archaeon DNA polymerases.

Three elements combine to enhance chain terminator incorporation

5

In summary, three elements are identified to enhance chain terminator incorporation. First, use of specific dye adducts attached to the nucleoside base. Second, use of acyclo base analogs. Third, use of archaeon DNA polymerase variants with increased ability to incorporate nucleotides with modified sugars. Each of the three elements alone enhances the desired incorporation. To a first approximation, these elements appear to act in an additive way, with the combined effects of two elements being greater than either alone, and the combined effects of all three elements being greater than any two elements alone.

15

While dye derivatives are emphasized in this application, the skilled worker will also recognize that other types of modified nucleotides could also be used. For example, the fluorescein moiety can also act as a hapten in antibody-based detection systems. Similarly, other nucleotide modifications that cross-react with a second molecule that can act in a detection scheme will also function in this invention.

20

While this invention treats incorporation of various terminators into otherwise native DNA on a DNA template, one skilled in the art will recognize that either the template or primer utilized in the reaction may contain nucleotide analogs that allow them to be the functional equivalent of such substrates. Such analogs might include, but not be limited to, thiophosphate backbone linkages, substituted bases and ribonucleotides. The invention requires only that the DNA polymerase employed be able to direct incorporation of the terminator in a base-specific fashion.

25

30

One significant advantage arising from more efficient incorporation of dye-terminators is a reduction in the amount of dye-terminator needed in the polymerization reaction. As a consequence, lower backgrounds and

increased sensitivity of detection are anticipated due to the higher ratio of incorporated to unincorporated substrate.

Applications of the invention

Recognition of the ability of this class of DNA polymerase and polymerase variants to incorporate dye-terminators has a broad range of applications. One obvious application is in DNA sequencing, where incorporation of dye-terminators forms the basis of numerous automated sequencing technologies (Lee, *et al.*, *Nucleic Acids Res.* **20**:2471-2483 (1992); Example 13, Figure 10). Past experience has shown nucleotide incorporation varies depending on the sequence context, and that this variability is specific for different DNA polymerases (Lee, *supra.* (1992); Brandis, *Nucleic Acids Res.* **27**:1912-1918 (1999); Parker, *Biotechniques* **19**:116-211 (1995)). It can be reasonably expected that some regions that are difficult to sequence with current protocols could more easily be rendered with the combinations of polymerases and terminators disclosed here. Since the insertion properties of these polymerases are different from those of Taq and related DNA polymerases, it may also be possible to mix archaeon DNA polymerases with other polymerases in sequencing reactions to allow more uniform signal incorporation, with both enzymes contributing to the final sequencing product. This mixing could be extended to other applications where dye-substituted bases are incorporated.

Other related applications such as fluorescence dideoxy finger-printing (F-ddF: Ellison, *et al.*, *Biotechniques* **17**:742-753 (1994)) can make use of this invention. Such applications require only short-range DNA sequence determinations, some as short as a single base. Discovery of compatible combinations of archaeon DNA polymerases and dye terminators allows such determinations to go forward, as might be required in detecting single nucleotide polymorphisms (SNPs; US Patent Nos. 5,888,819, 5,952,174, 6,004,744, and 6,013,43). For example, allele-specific primers can be extended by dye-labeled terminators, and the specific nucleotide inserted later detected by either fluorescence polarization (Chen, *et al.*, *Genome*

Research **9**:492-498 (1999)) or by fluorescence resonance energy transfer (Chen and Kwok, *Nucleic Acids Res.* **25**:347-353 (1997)).

5 The ability to insert non-standard nucleotides is also useful in sequencing applications employing mass spectroscopy. One limitation of multiplex genotyping by mass spectrometry is distinguishing the masses of oligonucleotide primers extended by a single nucleotide. By increasing the difference in mass between the four nucleotides added, increased resolution could be achieved, allowing analysis of larger oligomers, and increased
10 confidence in multiplex analysis where a large number of different molecular weights will need to be determined (Ross, *et al.*, *Nature Biotechnology* **16**:1347-1351 (1998)). Of course, incorporation of acyclo-derivatives without dyes could also be employed in this application.

15 The present invention is further illustrated by the following Examples. These Examples are provided to aid in the understanding of the invention and are not construed as limitations thereof.

20 The references cited above and below are herein incorporated by reference.

EXAMPLE 1

A titration assay to measure the relative efficiency of modified nucleotide incorporation

25

 The relative efficiency of modified nucleotide incorporation was assessed using variations of the assay described by Gardner and Jack (*supra.*). A primed single-stranded DNA substrate is incubated in a reaction
30 mixture containing a fixed concentration of dNTPs and increasing amounts of the modified nucleotide. Reactions can either be isothermal, or can be linearly amplified by thermal cycling using stages of denaturation, annealing and primer extension. Following the reaction, terminated extension products are separated by denaturing polyacrylamide gel electrophoresis, and the
35 separated products detected either by virtue of labels attached to the primer

-28-

(*e.g.*, 5'-[³²P] end-labeled) or terminator (*e.g.*, dye-labels) using methods commonly known in the art, such as autoradiography and fluorescent scanning.

Once the spectrum of termination products are determined, a comparison of the length, uniformity and clarity of this pattern is used to evaluate the relative incorporation of the modified nucleotide. The more readily incorporated terminators will produce a given pattern when present in lower concentrations than the comparison standard. Alternatively, the banding pattern at a given concentration of modified nucleotide can be compared between two or several compounds. The compounds producing shorter termination products at a given concentration are those that are more efficiently incorporated by the DNA polymerase. This latter method can theoretically be performed using a single analog concentration, although it is more desirable to use multiple concentrations to provide greater opportunities for comparison.

A control reaction, containing no terminator, confirms that the polymerase is able to fully extend the primer (approximately 7200 bp in the case of M13mp18) in the absence of the terminator. Thus, the bands observed in other reactions arise from terminator incorporation rather than incomplete replication by the DNA polymerase.

EXAMPLE 2

Dye-ddCTP derivatives differ in incorporation efficiency

A variety of available dye-labeled ddCTP derivatives (Table 2) were analyzed and compared to test for incorporation by Vent® (exo-) DNA polymerase. Primed M13mp18 substrate was formed as previously described (Kong, *et al.*, *supra.*). As in all the examples, all reaction components were from New England Biolabs (Beverly, MA), except where indicated. Incorporation of modified bases was assayed by mixing 2.5 μ l of 2X reaction cocktail (0.04 μ M 5' [³²P] end-labeled #1224-primed M13mp18, 2X ThermoPol Buffer, 0.04 U/ μ l thermostable pyrophosphates, 80 μ M dNTP, 0.15 U/ μ l DNA polymerase) with 2.5 μ l of nucleotide analog solution to yield the final ratios of analog : dCTP indicated in the figures. After

-29-

incubating at 72°C for 15 minutes, the reactions were halted by the addition of 4 μ l CircumVent® stop/dye (0.3% xylene cyanole FF., 0.3% bromophenol blue, 0.37% EDTA, pH 7.0). Samples were then heated at 72°C for 3 minutes and separated on a QuickPoint DNA sequencing gel (NOVEX, San Diego, CA) run at 1200 volts. The gel was fixed by soaking in 10% acetic acid/10% ethanol, dried, and polymerization products visualized by autoradiography. Examples of these reactions are given in Figure 1.

The extent of dye-terminator was determined by visual inspection of the autogradiograms. Evaluations followed the outlines given in Example 1, and are recorded in Table 2.

Table 2

Commercially available dye-terminators from NEN Life Science Products Catalog

<u>Dye-Terminator</u>	<u>Catalog Number</u>	<u>Incorporation</u>
JOE-ddATP	NEL486	less
JOE-ddCTP	NEL485	
JOE-ddGTP	NEL487	
JOE-ddUTP	NEL484	
TAMRA-ddATP	NEL474	better
TAMRA-ddCTP	NEL473	
TAMRA-ddGTP	NEL475	
TAMRA-ddUTP	NEL472	
FAM-ddATP	NEL482	less
FAM-ddCTP	NEL481	
FAM-ddGTP	NEL483	
FAM-ddUTP	NEL480	
ROX-ddATP	NEL478	better
ROX-ddCTP	NEL477	
ROX-ddGTP	NEL486	
ROX-ddUTP	NEL479	
Fluorescein-12-ddATP	NEL402	
Fluorescein-12-ddCTP	NEL400	
Fluorescein-12-ddGTP	NEL403	
Fluorescein-12-ddUTP	NEL401	
R6G-ddCTP	NEL489	better slightly worse
R110-ddCTP	NEL493	

Table 2 (continued)

Non-commercially available dye-terminators tested***Dye-Terminator***

5	ROX-acyclo-CTP	better
	TAMRA-acyclo-CTP	better
	FI-12-acyclo-CTP	better
	IRD40-ddCTP	similar
	IRD700-ddCTP	slightly better
10	IRD700-acyclo-NTP*	better
	Cyanine 3-ddCTP	less
	Cyanine 5-ddCTP	less
	BODIPY® TR-ddCTP	better
	BODIPY®TMR-ddCTP	better
15	BODIPY®R6G-ddCTP	similar
	BODIPY®FI-ddCTP	similar
	BODIPY®FL-acyclo-GTP	better
	IRD40-acyclo-NTP*	better
20	R6G-acyclo-ATP	better

Analogues labeled with * were tested for all four bases. Incorporation refers to incorporation relative to ddNTP. These alkynylamino-acyclic analogs are covered by U.S. Patent Nos. 5,047,519 and 5,151,507 issued to NEN Life Science Products, Inc.

EXAMPLE 3***Blast comparison of Family B DNA polymerases***

One method of classifying and categorizing proteins is by primary amino acid sequence alignment. It is generally accepted that high degrees of primary sequence similarity suggest similar function, and thus can be predictive of physical and enzymatic properties common between the compared proteins. A sampling of archaeon DNA polymerases, along with representatives of other Family B and Family A DNA polymerases, were compared using the sequence alignment program Blastp. This program searches for a maximal collinear sequence alignment between test

sequences, with output reported in terms of sequence identity, sequence similarity (where paired amino acid residues have similar characteristics), and gaps introduced to maintain the alignment.

5 The source for sequence information was the ncbi server at the internet site: <http://www.ncbi.nlm.nih.gov>, and accession numbers derived from that site are listed along with the source organism in Table 3. Blastp comparisons were run pairwise with either the Vent® DNA polymerase amino acid sequence, with the following program parameters:

10 matrix: 0BLOSUM62

 gap open: 11

 gap extension: 1

 x_dropoff: 50

 expect: 10

15 wordsize: 3

 filter: off

Comparisons were done *via* the internet site:

<http://www.ncbi.nlm.nih.gov/gorf/bl2.html>. Column four of Table 3 report % identity/% positives/% gaps for such comparisons. Entries marked “none”
20 returned “no significant similarity found” on blastp analysis.

 The polymerases can be assigned to several groups based on this analysis. First, those polymerases with greater than about 70% sequence identity with Vent® DNA polymerase. In the sequences tested, such
25 enzymes were derived from *Thermococcus* or *Pyrococcus* species, although examples from other species may also be found. Second, an intermediate group of between about 30 and 70% identity, with examples listed here deriving from *Pyrodictium* and *Methanococcus* species. Third, Family B DNA polymerases with less than about 30% identity, including both archaeon, viral
30 and eukaryotic DNA polymerases. Fourth, Family A DNA polymerases with no significant similarity detected by the analysis.

 Those DNA polymerases with higher identity percentages are considered to be more likely to share base analog incorporation properties
35 with Vent® DNA polymerase. The archaeon DNA polymerases reported in

later examples derive from the first group, consistent with this prediction. The next most likely group to contain similar sequence characteristics are those with about 30-70% sequence identity. Similarly, the Family A DNA polymerases are shown to have very different properties, as might be suggested by the lack of sequence identity detected by this sequence analysis. The intermediate groups, including the second and third groups above, would be expected to behave more like Vent® DNA polymerase than like the Family A DNA polymerases.

Table 3 also lists an outgrowth of the sequence comparisons, namely an alignment of the conserved region mutagenized in several of the polymerases (see also Table 1). Notable is the conservation of the key, underlined residues in polymerases with high sequence identity to Vent® DNA polymerase, encouraging the view that homologous mutations in those related polymerases will have similar effects on nucleotide incorporation as observed in Vent® DNA polymerase.

EXAMPLE 4

Purification of 9°N™ (exo-) DNA polymerase

The 9°N™ (exo-) DNA polymerase (referred to as the "AIA" mutant) was constructed, grown and expressed in a T7 expression system as described in Southworth, *et al.* (*supra.*). Purification followed the general outline described in that reference and, except where noted, was at 0-4°C. The cell pellet (380 g) was suspended in 1.14 liter of Buffer A (20 mM KPO₄ (pH 6.8), 0.1 mM EDTA, 0.05% Triton X-100, 0.1 M NaCl, 10% glycerol). Cells were lysed by multiple passages through a Manton Gaulin homogenizer, using a cooling coil to keep the homogenate temperature below 20°C. The extract was clarified by centrifugation for 40 min. in a Sharples Type 16 centrifuge at 15,000 rpm (Fraction I, volume 1.5 liters).

The cleared extract was heated to 75°C for 10 minutes, and then cooled on ice. Insoluble material was removed by centrifugation for 35 min. at 4 krpm in a Beckman JS-4.2 rotor (Fraction II, 0.85 liters).

-33-

The cleared extract was heated to 75°C for 10 minutes, and then cooled on ice. Insoluble material was removed by centrifugation for 35 min. at 4 krpm in a Beckman JS-4.2 rotor (Fraction II, 0.85 liters).

5 Fraction II was passed through a 0.7 liter (9.5x10 cm) DEAE-cellulose column, equilibrated in Buffer A containing 1 mM DTT and immediately applied to a 235 ml (5x12 cm) phosphocellulose column equilibrated in the same buffer. The latter column was washed with 0.5 liter of buffer A containing 1 mM DTT, and eluted with a 2 liter linear gradient of NaCl (0.1-1.0 M).
10 Polymerase activity was assayed, and peak fractions pooled (Fraction III, volume 0.4 liter, approximately 0.8 g protein).

15 Fraction III was dialyzed against buffer B (20 mM TrisHCl (pH 7.6), 0.1 M NaCl, 1 mM DTT, 0.1 mM EDTA, 10% glycerol), and passed through a 49 ml (2.5x10 cm) DEAE-cellulose column. The column was washed with 50 ml of buffer B, and the wash was combined with the flow-through fractions (Fraction IV, volume 0.45 liter).

20 Fraction IV was dialyzed against buffer C (20 mM TrisHCl (pH 7.5), 0.05 M NaCl, 1 mM DTT, 0.1 mM EDTA, 10% glycerol). Solid $(\text{NH}_4)_2\text{SO}_4$ was added (101 g) to a final concentration of 1.7 M, and the solution was filtered through a 0.22 μm filter. The filtered solution was applied to a 0.05 liter phenyl-sepharose column (2.5x10 cm) equilibrated in buffer C containing 1.7 M $(\text{NH}_4)_2\text{SO}_4$. The column was washed with 0.1 liter of buffer C containing
25 1.7 M $(\text{NH}_4)_2\text{SO}_4$, and eluted with a 0.5 liter linear gradient of $(\text{NH}_4)_2\text{SO}_4$ (1.7-0 M) in buffer C. Column fractions were assayed for polymerase activity and fractions containing the peak of polymerase activity were pooled (Fraction V, 90 ml, approximately 0.3 g protein).

30 Fraction V was dialyzed against buffer B, and loaded onto a 53 ml (2.5x10 cm) Affigel Blue column equilibrated in the same buffer. Following loading, the column was washed with 0.1 liter of buffer B, and eluted with a 0.5 liter linear gradient of NaCl (0.1-1.35 M) in buffer B. Fractions were assayed for polymerase activity, and the peak of the activity fractions were
35 pooled and dialyzed into storage buffer (10 mM TrisHCl (pH 7.4), 0.1 M

EXAMPLE 5

Dye-acyclo-CTP derivatives are more efficiently incorporated than dye-ddCTP derivatives by Vent® (exo-) DNA polymerase

Acyclo-NTPs, similar to ddNTPs, lack a free 3-OH termini, and are expected to act as chain terminators in DNA polymerase reactions. The ability of acyclo-NTPs with dye-derivatized bases to act as chain terminators was tested using a titration assay. Incorporation of ROX-ddCTP, TAMRA-ddCTP and IRD700-ddCTP were compared to that of ROX-acyclo-CTP, TAMRA-acyclo-CTP and IRD700-acyclo-CTP, respectively, using Thermo Sequenase™ and Vent® (exo-) DNA polymerases.

Incorporation of modified bases was assayed by mixing 2.5 μ l of 2X reaction cocktail (0.04 μ M 5' [³²P] end-labeled #1224-primed M13mp18, 2X Thermopol Buffer, 0.04 U/ μ l thermostable pyrophosphates, 80 μ M dNTP, 0.15 U/ μ l DNA polymerase) with 2.5 μ l of nucleotide analog solution to yield the final ratios of analog : dCTP indicated in the figures. After incubating at 72°C for 15 minutes, the reactions were halted by the addition of 4 μ l CircumVent® stop/dye (0.3% xylene cyanole FF., 0.3% bromophenol blue, 0.37% EDTA (pH 7.0)). Samples were then heated at 72°C for 3 minutes and separated on a QuickPoint DNA sequencing gel (NOVEX, San Diego, CA) run at 1200 volts. The gel was fixed by soaking in 10% acetic acid/10% ethanol, dried, and polymerization products visualized by autoradiography. Examples of these reactions are given in Figure 2.

A comparison of banding patterns revealed shorter terminator products for the acyclo-CTP, as opposed to the ddCTP, derivatives when using Vent® (exo-) DNA polymerase. This was the case for ROX, TAMRA and IRD700 derivatives. Thus, the acyclo derivatives are more efficiently incorporated than their dideoxy equivalents. In contrast, Thermo Sequenase™ showed a preference for dideoxy derivatives, as evident in the collection of shorter termination products for those ROX, TAMRA and IRD700 derivatives.

EXAMPLE 6

Sensitivity to dye-acyclo-CTP terminators is shared by a variety of archaeon DNA polymerases

The sequence similarity among archaeon Family B DNA polymerases raises the possibility that they will function similar to Vent® (exo-) DNA polymerase with respects to incorporation of dye terminators. This proposition was tested using ROX-acyclo-CTP in a titration assay to compare the performance of Vent® (exo-), Deep Vent® (exo-), Pfu (exo-) and 9°N™ (exo-) DNA polymerases. A titration assay of the type described in Example 1 was used.

Briefly, a 2X reaction cocktail was prepared on ice containing 0.06 μ M 5'-[³²P] #1224 primed single-stranded M13mp18, 2X ThermoPol Buffer (20 mM KCl, 40 mM Tris-HCl (pH 8.8 at 25°C), 20 mM (NH₄)₂SO₄, 4 mM MgSO₄, 0.2% Triton X-100), 0.04 U/ μ l thermostable inorganic pyrophosphatase and 80 μ M dNTP. The 2X cocktail was split into aliquots and Vent® (exo-), Deep Vent® (exo-) or 9°N™ (exo-) DNA polymerase was added to a final concentration of 0.06 U/ μ l. Another 2X cocktail was made with conditions recommended by the manufacturer (Stratagene, La Jolla, CA) for Pfu (exo-) DNA polymerase containing 0.06 μ M 5'-[³²P] #1224 primed single-stranded M13mp18, 2X Pfu Buffer (20 mM KCl, 20 mM (NH₄)₂SO₄, 40 mM Tris-HCl (pH 8.75), 4 mM MgSO₄, 0.2% Triton X-100, 0.2 mg/ml BSA), 0.04 U/ μ l thermostable inorganic pyrophosphatase, and 80 μ M dNTP to which Pfu (exo-) DNA polymerase was added to a final concentration of 0.06 U/ μ l. A 2.5 μ l aliquot of 2X reaction cocktail was mixed with 2.5 μ l of nucleotide analog to yield the final ratios of analog : dCTP indicated in the figures. Control extensions added 2.5 μ l of dH₂O to 2.5 μ l of reaction mix, and demonstrated that polymerization proceeded without termination in the absence of ROX-acyclo-CTP. Following mixing, reactions were immediately incubated at 72°C for 20 minutes.

Reactions were stopped by the addition of 4 μ l NEB Stop/Loading Dye Solution (deionized formamide containing: 0.3% xylene cyanole FF,

-36-

0.3% bromophenol blue, 0.37% EDTA (pH 7.0)) and heated at 72°C for 3 minutes. A 1 μ l aliquot was loaded onto a QuickPoint (NOVEX) mini-sequencing gel and run at 1200 V for 10 minutes. The gel was fixed, washed, and dried according to manufacturer's instructions and polymerization products visualized by autoradiography (Figure 3).

Based on analysis of termination fragment lengths, ROX-acyclo-CTP was incorporated by all four archaeon DNA polymerases Vent® (exo-), Deep Vent® (exo-), Pfu (exo-) and 9°N™ (exo-). At high concentrations (2 : 1 molar ratio) of ROX-acyclo-CTP : dCTP, termination products were very short due to the efficient incorporation of the terminator. The similar range and quality of termination fragments suggesting that all four archaeon DNA polymerases incorporate ROX-acyclo-CTP with comparable efficiency.

EXAMPLE 7

Incorporation of dye-labeled terminators is enhanced by Vent® (exo-)/A488L DNA polymerase

The ability of Vent® (exo-)/A488L DNA polymerase to incorporate dye-labeled terminators was evaluated using the titration assay described in Example 1. A variety of available dye-labeled ddCTP derivatives (Table 2) were analyzed and compared to test for incorporation by Vent® (exo-) DNA polymerase. Primed M13mp18 substrate was formed as previously described (Kong, *et al.*, *supra.*). As in all the examples, all reaction components were from New England Biolabs (Beverly, MA), except where indicated. Incorporation of modified bases was assayed by mixing 2.5 μ l of 2X reaction cocktail (0.04 μ M 5' [³²P] end-labeled #1224-primed M13mp18, 2X Thermopol Buffer, 0.04 U/ μ l thermostable inorganic pyrophosphatase, 80 μ M dNTP, 0.15 U/ μ l DNA polymerase) with 2.5 μ l of nucleotide analog solution to yield the final ratios of analog : dCTP indicated in the figures. After incubating at 72°C for 15 minutes, the reactions were halted by the addition of 4 μ l CircumVent® stop/dye (0.3% xylene cyanole FF., 0.3% bromophenol blue, 0.37% EDTA, pH 7.0). Samples were then heated at 72°C for 3 minutes and separated on a QuickPoint DNA sequencing gel (NOVEX, San Diego, CA) run at 1200 volts. The gel was fixed by soaking

-37-

in 10% acetic acid/10% ethanol, dried, and polymerization products visualized by autoradiography. Examples of these reactions are given in Figure 4.

In each case dye-terminators were incorporated more efficiently by Vent® (exo-)/A488L DNA polymerase than by the parent Vent® (exo-) DNA polymerase. Despite this increase, the relative efficiency of incorporation among the various dye-terminators was the same for both enzymes.

EXAMPLE 8

Alternate DNA polymerase variants can also enhance the incorporation of terminators

Additional Vent® (exo-) DNA polymerase variants were also tested for their ability to enhance incorporation of dye-terminators. The variant Y499L was compared with the parental (exo-) polymerase, along with the A488L variant in its ability to incorporate ROX-ddCTP.

As in previous examples, the efficiency of analog incorporation was determined using a titration assay, using varying concentrations of terminators. Briefly, a 2X reaction cocktail was prepared on ice containing 0.04 μ M single-stranded M13mp18 primed with 5'-[³²P] end-labeled #1224 primer, 2X ThermoPol Buffer (20 mM KCl, 40 mM Tris-HCl (pH 8.8 at 25°C), 20 mM (NH₄)₂SO₄, 4 mM MgSO₄, 0.2% Triton X-100), 0.04 U/ μ l thermostable inorganic pyrophosphatase and 80 μ M dNTP. The 2X cocktail was split into aliquots and Vent® (exo-), Vent® (exo-)/A488L or Vent® (exo-)/Y499L DNA polymerase was added to a final concentration of 0.06 U/ μ l. A 2.5 μ l aliquot of this 2X reaction cocktail was mixed with 2.5 μ l of a nucleotide analog mixture, resulting in the final ratios of analog : dCTP indicated in the figures. Control reactions mixed 2X reaction cocktail with an equal volume of dH₂O. Reactions were immediately incubated at 72°C for 15 minutes.

Reactions were stopped by the addition of 4 μ l NEB Stop/Loading Dye Solution (deionized formamide containing: 0.3% xylene cyanole FF,

-38-

0.3% bromophenol blue, 0.37% EDTA (pH 7.0)) and heated at 72°C for 3 minutes. A 1 μ l aliquot was loaded onto a QuickPoint (NOVEX) mini-sequencing gel and run at 1200 V for 10 minutes. The gel was then fixed, washed, and dried according to manufacturer's instructions and reaction products were visualized by autoradiography (Figure 5).

Both the A488L and Y499L variants of Vent® (exo-) DNA polymerase were better able to incorporate ROX-ddCTP than the parent Vent® (exo-) DNA polymerase (Figure 5) as evidenced in the shorter termination products produced by those variants at the same analog concentrations. Incorporation by the two variants was comparable, with slightly more efficient terminator incorporation by the A488L variant. Approximately 5-10-fold lower concentrations of ROX-ddCTP were required to produce equivalent banding patterns with the variants compared to the parent enzyme. These enzymes are thus useful tools for greater incorporation of chain terminating nucleotides.

EXAMPLE 9

Generation of 9°N™ DNA polymerase variants

Production and purification of Vent® DNA polymerase variants was as described (Gardner and Jack, *supra*). This led to enzyme preparations that were substantially purified, meaning separated from contaminants affecting the performance of the enzyme, such as contaminating exo- and endonucleases, alternate polymerases and endogenous nucleotides. Purification of 9°N™ (exo-) DNA polymerase and the A485L variant of that enzyme used the same protocols.

An expression vector for the A485L variant of 9°N™ DNA polymerase was created using PCR mutagenesis (Colosimo, *et al. Biotechniques* **26**:870-873 (1999)) of the expression construct pNEB917, a derivative of pNEB915 encoding an exonuclease-deficient (AIA) form of the polymerase (Southworth, *et. al, Proc. Natl. Acad. Sci. USA* **93**:5281-5285 (1996)).

The mutagenesis used two successive PCR reactions. The first stage reactions (0.05 ml) contained 1X Thermopol buffer (New England Biolabs, Beverly, MA), 50 ng/ml pNEB915 template DNA, 0.25 mM dNTPs, 0.5 μ M oligonucleotide #216-153 (SEQ ID NO:1; Table 4), 0.5 μ M oligonucleotide #175-70 (SEQ ID NO:2; Table 4), 0.1 mg/ml bovine serum albumen and 2 mM added MgSO_4 in a 0.2 ml thin-wall PCR tubes. One unit of Vent® DNA polymerase was added to the reaction mixture, and the tube containing the mixture was heated at 94°C for three minutes, followed by 25 cycles of 94°C (15 seconds), 58°C (15 seconds), 72°C (60 seconds). Evaluation of the reaction product on an agarose gel revealed a band of the expected molecular weight.

Table 4**Oligonucleotide Sequences**

<u>Oligo #</u>	<u>Sequence</u>
216-153 (SEQ ID NO:1)	CAGGCAGAGGCTTATAAAAATCCTCGCCAACAGCTT
175-70	GGTGGCAGCAGCCAACCTCAGCTTCCT (SEQ ID NO:2)
216-155	GATTCTCATGATAAGCTACGCCGA (SEQ ID NO:3)

The second round of PCR was accomplished by diluting the above PCR sample either 250- or 500-fold into 0.1 ml reaction mixtures containing: 1X Thermopol buffer, 0.1 mg/ml bovine serum albumen, 0.25 mM dNTPs, 0.5 μ M oligonucleotide #175-70 (SEQ ID NO:2; Table 4), 0.5 μ M oligonucleotide #216-155 (SEQ ID NO:3; Table 4), and 0, 2, 4, 6 or 8 mM added MgSO_4 , again in 0.2 ml thin-walled PCR tubes. After addition of one unit of Vent® DNA polymerase to each reaction mixture, the sample was heated at 94°C for three minutes, followed by 25 cycles of 94°C (15 seconds), 58°C (15 seconds), 72°C (90 seconds). Aliquots of each reaction were analyzed by agarose gel electrophoresis and found to contain a band

-40-

of the expected size (about 1.5 kb). Samples containing 2-8 mM added MgSO_4 were pooled, phenol extracted and ethanol precipitated.

5 The precipitated sample was suspended in 0.1 ml of 1X NEBuffer 2, and cut sequentially with the restriction endonucleases BamHI (100 units for 1 hour at 37°C) and BsiWI (75 units for 1 hour at 55°C). The plasmid pNEB917 was similarly digested with the same enzymes. The reaction products from both samples were separated on a 0.7% agarose gel in TBE buffer containing 0.5 $\mu\text{g/ml}$ ethidium bromide. The prominent approximately 10 1.5 kb band derived from the PCR sample and the approximately 7 kb band derived from pNEB917 were excised and eluted using an Elutrap apparatus in 0.5X TBE, using conditions specified by the manufacturer (Schliecher & Schuell, Keene, NH). The eluted DNAs were phenol extracted and ethanol precipitated. After suspension of the DNA pellet in TE buffer, the 15 samples were quantified by running small aliquots on an agarose gel, and comparing the samples with molecular mass and weight standards.

The eluted fragments were ligated, and ampicillin resistant transformants were selected and screened by cleavage with PstI, a site not 20 present on pNEB917, but which would be gained if the mutagenesis was successful. One construct displaying the PstI site was named pEAC3, and was used for expression of 9°N™ (exo-/A485L) DNA polymerase.

25 Expression and purification of the variant DNA polymerase was as described for Vent® (exo-)/A488L (Gardner and Jack, *supra*).

EXAMPLE 10

30 ***Vent® (exo-)/A488L and 9°N™ (exo-)/A485L DNA polymerases both efficiently incorporate ROX-ddCTP***

The high degree of sequence identity in archaeon DNA polymerases suggests that variants similar to the A488L variant described in Example 7 should function similarly with respects to dye-terminator incorporation. 35 Accordingly, Vent® (exo-)/A488L and 9°N™ (exo-)/A485L DNA

polymerases were compared in their ability to incorporate both ddCTP and ROX-ddCTP.

A 2X reaction cocktail was prepared on ice containing 0.04 μ M single-stranded M13mp18 primed with 5'-[32 P] end-labeled #1224 primer, 2X ThermoPol Buffer (20 mM KCl, 40 mM Tris-HCl (pH 8.8 at 25°C), 20 mM (NH₄)₂SO₄, 4 mM MgSO₄, 0.2% Triton X-100), 0.04 U/ μ l thermostable inorganic pyrophosphatase and 80 μ M dNTP. The 2X cocktail was split into aliquots and Vent® (exo-), Vent® (exo-)/A488L or 9°N™ (exo-)/A485L DNA polymerase was added to a final concentration of 0.04 U/ μ l. A 2.5 μ l aliquot of this 2X reaction cocktail was mixed with 2.5 μ l of a nucleotide analog mixture, resulting in the final ratios of analog : dCTP indicated in the figures. Control reactions mixed 2X reaction cocktail with an equal volume of dH₂O. Reactions were immediately incubated at 72°C for 15 minutes.

Reactions were stopped by the addition of 4 μ l NEB Stop/Loading Dye Solution (deionized formamide containing: 0.3% xylene cyanole FF, 0.3% bromophenol blue, 0.37% EDTA (pH 7.0)) and heated at 72°C for 3 minutes. A 1 μ l aliquot was loaded onto a QuickPoint (NOVEX) mini-sequencing gel and run at 1200 V for 10 minutes. The gel was then fixed, washed, and dried according to manufacturer's instructions and reaction products were visualized by autoradiography (Figure 6).

Reactions of Vent® DNA polymerase (exo-) and 9°N™ (exo-) with ddCTP were characterized by a faint series of high molecular weight bands, reflecting relatively weak incorporation of ddCTP under these conditions. The same pattern of termination products is observed when ROX-ddCTP is used at approximately 25-fold lower concentrations, a measure of the increased efficiency of incorporation of the ROX analog by this enzyme.

The 9N™ (exo-)/A485L DNA polymerase variant mimicked the enhanced incorporation noted for the Vent® (exo-)/A488L DNA polymerase both with respects to relative incorporation of ddCTP and ROX-ddCTP (Figure 6). With each enzyme depicted in Figure 6, ROX-ddCTP is incorporated about 10-fold better than ddCTP (compare 10 : 1 vs. 1 : 1

lanes for ddCTP : dCTP vs. ROX-ddCTP : dCTP, respectively). Additional comparison of the parental and variant enzymes showed a further approximately 10-fold enhancement of either ROX-ddCTP or ddCTP incorporation.

5

EXAMPLE 11

Archaeon DNA polymerase variants display enhanced incorporation of dye-acyclo-NTPs relative to dye-ddNTPs

10

15

While the previous examples illustrate the utility of using archaeon DNA polymerase variants to enhance incorporation of dye-ddNTP terminators, their utility in incorporation of dye-acyclo-NTPs was unknown. The experiments in this example illustrate that terminator incorporation is further enhanced when archaeon variants are used in combination with acyclo terminators. Two sets of experiments established these facts.

20

25

In the first experiment, a 2X reaction cocktail was prepared on ice containing 0.04 μ M single-stranded M13mp18 primed with 5'-[³²P] end-labeled #1224 primer, 2X ThermoPol Buffer (20 mM KCl, 40 mM Tris-HCl (pH 8.8 at 25°C), 20 mM (NH₄)₂SO₄, 4 mM MgSO₄, 0.2% Triton X-100), 0.04 U/ μ l thermostable inorganic pyrophosphatase and 0.08 mM dNTP. The 2X cocktail was split into aliquots and Vent® (exo-)/A488L DNA polymerase was added to a final concentration of 0.04 U/ μ l (0.08 U/ μ l for IRD700-acyclo-CTP). A 2.5 μ l aliquot of this 2X reaction cocktail was mixed with 2.5 μ l of a nucleotide analog mixture, resulting in the final ratios of analog : dCTP indicated in Figure 7. Reactions were immediately incubated at 72°C for 20 minutes.

30

Reactions were stopped by the addition of 4 μ l NEB Stop/Loading Dye Solution (deionized formamide containing: 0.3% xylene cyanole FF, 0.3% bromophenol blue, 0.37% EDTA (pH 7.0)) and heated at 72°C for 3 minutes. A 1 μ l aliquot was loaded onto a QuickPoint (NOVEX) mini-sequencing gel and run at 1200 V for 10 minutes. The gel was then fixed,

-43-

washed, and dried according to manufacturer's instructions and reaction products were visualized by autoradiography (Figure 7).

As noted with the parental Vent® (exo-) DNA polymerase, incorporation of dye-acyclo-CTP was enhanced relative to dye-ddCTP for IRD700, ROX and TAMRA analogs. Thus, this variant retains the favorable acyclo-NTP incorporation characteristics of the parent.

In a related experiment dye-acyclo-CTP incorporation by the analogous 9°N™ (exo-)/A485L variant was evaluated. A 2X reaction cocktail was prepared on ice containing 0.1 mg/ml single-stranded M13mp18, 0.1 μM 5' [³²P] labeled primer #1224, 2X ThermoPol Buffer (20 mM KCl, 40 mM Tris-HCl (pH 8.8 at 25°C), 20 mM (NH₄)₂SO₄, 4 mM MgSO₄, 0.2% Triton X-100), 0.04 U/μl thermostable inorganic pyrophosphatase and 80 μM dNTP. The 2X cocktail was split in half and Vent® (exo-)/A488L, 9°N™ (exo-)/A485L or Thermo Sequenase™ DNA polymerase was added to a final concentration of 0.06 U/μl. A 2.5 μl aliquot of 2X reaction cocktail was mixed with 2.5 μl of an 80μM nucleotide analog mix in 0.5 ml tubes and immediately incubated at 72°C for 20 minutes.

Reactions were stopped by the addition of 4 μl NEB Stop/Loading Dye Solution (deionized formamide containing: 0.3% xylene cyanole FF, 0.3% bromophenol blue, 0.37% EDTA (pH 7.0)) and heated at 72°C for 3 minutes. A 1 μl aliquot was loaded onto a QuickPoint (NOVEX) mini-sequencing gel and run at 1200 V for 10 minutes. The gel was then fixed, washed, and dried according to manufacturer's instructions and reaction products were visualized by autoradiography (Figure 8).

Vent® and 9°N™ DNA polymerase variants displayed comparable incorporation of all dye-acyclo-CTP analogs tested, establishing the interchangeability of these analogous variants for this invention.

EXAMPLE 12***Acyclo-GTP is more efficiently incorporated than ddGTP by archaeon DNA polymerases***

5 Example 5 illustrated the increased efficiency of dye-acyclo-NTPs compared to the corresponding dye-ddNTPs. While these results strongly suggest that the increase in incorporation efficiency arises from the acyclo modification, a direct test was performed. The ability of both Thermo Sequenase™ and 9°N™ (exo-)/A485L to incorporate acyclo-GTP and
10 ddGTP was evaluated using the titration assay.

A 2X reaction cocktail was prepared on ice containing 0.1 mg/ml single-stranded M13mp18, 0.1 μM 5'-[³³P] end-labeled primer #1224, 2X
15 ThermoPol Buffer (20 mM KCl, 40 mM Tris-HCl (pH 8.8 at 25°C), 20 mM (NH₄)₂SO₄, 4 mM MgSO₄, 0.2% Triton X-100), 0.04 U/μl thermostable inorganic pyrophosphatase and 0.1 mM dNTPs. The 2X cocktail was split in half and 9°N™ (exo-)/A485L DNA polymerase or Thermo Sequenase™ was added to a final concentration of 0.04 U/μl. A 2.5 μl aliquot of 2X reaction
20 cocktail was mixed with 2.5 μl of a nucleotide analog mix to yield the final ratios of analog : dGTP indicated in the figures, and immediately placed in a thermal cycler preheated to 94°C. Reactions were thermal cycled as follows:

94°C 5 minutes

25 cycles at:

25 94°C 30 seconds

55°C 30 seconds

72°C 30 seconds

72°C 7 minutes

Reactions were stopped by the addition of 4 μl Stop/Loading Dye
30 Solution (deionized formamide containing: 0.3% xylene cyanole FF, 0.3% bromophenol blue, 0.37% EDTA (pH 7.0)) and heated at 72°C for 3 minutes. A 1 μl aliquot was loaded onto a QuickPoint (Novex) mini-sequencing gel and run at 1200 V for 10 minutes. The gel was then fixed, washed, and dried according to manufacturer's instructions and analyzed by
35 autoradiography.

-45-

The relative incorporation efficiency of ddGTP and acyclo-GTP by these two DNA polymerases was indicated by the analog concentration yielding equivalent banding patterns (Figure 9). For example, in a reaction using Thermo Sequenase™, 3:1 acyclo-GTP gave similar banding pattern to that seen with 1:9 ddGTP, indicating an approximate 27-fold preference for ddGTP over acyclo-GTP in these assays. On the other hand, 9°N™ (exo-)/A485L displayed similar banding patterns with 3:1 ddGTP and 1:3 μM acyclo-GTP, indicating an approximately 9-fold preference for acyclo-GTP over ddGTP in these assays.

EXAMPLE 13

DNA sequence analysis using archaeon DNA polymerase variants and dye-labeled acyclo-NTPs

The enhanced incorporation of modified nucleotides noted in the examples enables a new system of polymerases and reagents for use in automated DNA sequencing. These reactions rely upon incorporation of four chain terminators, each corresponding to one of the four bases normally present in DNA, and each labeled with a uniquely detectable fluorescent dye. The feasibility of such a reaction was tested using the following dye-labeled acyclo-NTPs: R6G-acATP (green), ROX-acCTP (red), BODIPY FL-acGTP (blue), and TAM-acUTP (yellow), where the color indicates the spectrum of the fluorescent emission and "ac" indicates acyclo derivatives. All analogs were obtained from NEN Life Science. Reaction products were separated by denaturing polyacrylamide gel electrophoresis, and detected via fluorescence upon laser excitation.

A reaction cocktail was prepared consisting of 50 ng/μl single-stranded M13mp18, 1 μM #1224 primer, 50 mM TrisHCl (pH 8.0 at room temperature), 8 mM MgSO₄, 0.2 M KCl, 0.1 mM dNTP, 0.1 μM R6G-acATP, 0.1 μM ROX-acCTP, 0.1 μM BODIPY® FL-acGTP, 0.25 μM TAM-acUTP, 0.02 U/μl thermostable inorganic pyrophosphatase and 0.04 U/μl 9°N™ (exo-)/A485L. Reactions were thermal cycled:

-46-

94°C 5 minutes

20 cycles of:

94°C 30 seconds

58°C 30 seconds

5 72°C 30 seconds

72°C 7 minutes

10 AmpliTaq® DNA polymerase, FS reactions were performed using materials acquired from and reaction conditions specified by the manufacturer (ABI PRISM™ Dye Terminator Cycle Sequencing Ready Reaction Kit protocol manual, P/N 402078 Revision A, August 1995, Perkin Elmer Corporation).

15 After the listed thermal cycling, unincorporated dye-labeled nucleotide terminators were separated by gel filtration using mini-columns (CentriSep, Princeton Separations), lyophilized and suspended in 5 µl formamide stop/dye (deionized formamide containing: 0.3% xylene cyanole FF, 0.3% bromophenol blue, 0.37% EDTA (pH 7.0)). Reactions were loaded onto a 4.75% urea gel, and reaction products were separated and detected by an 20 ABI377 automated DNA sequencer. DNA sequencing traces were processed and displayed using software Factura 2.0.1 and AutoAssembler 1.4.0 (Perkin-Elmer Corp.).

25 Termination fragments are detected by laser-excited fluorescent emission and plotted according to mobility, resulting in a pattern of peaks corresponding to each of the four dye terminators. The color of the peaks corresponds to the dye-acycloNTP that terminates the product. For example, a red peak on the trace would correspond to a product terminated by ROX-acCTP. Software assignment of peak identity appears above traces for both 30 AmpliTaq® DNA Polymerase, FS and 9°N™ (exo-)/A485L reactions, with the anticipated sequence appearing on the top line.

WHAT IS CLAIMED IS:

1. A method for site-specific incorporation of derivatized dideoxynucleotides into DNA comprising reacting an archaeon Family B DNA polymerase, a primed DNA template and nucleotide solution containing at least one derivatized dideoxynucleotide to produce fragments of DNA with the derivatized dideoxynucleoside covalently attached to the 3' terminal residue wherein the derivatized dideoxynucleotide is incorporated more efficiently than the corresponding underivatized dideoxynucleotide.

2. A method for site-specific incorporation of acyclonucleotides into DNA comprising reacting an archaeon Family B DNA polymerase, a primed DNA template and nucleotide solution containing at least one acyclonucleotide to produce fragments of DNA with the acyclonucleotide covalently attached to the 3' terminal residue.

3. A method for site-specific incorporation of derivatized acyclonucleotides into DNA comprising reacting an archaeon Family B DNA polymerase, a primed DNA template and nucleotide solution containing at least one derivatized acyclonucleotide to produce fragments of DNA with the derivatized acyclonucleotide covalently attached to the 3' terminal residue.

4. The method of claims 1 or 3 where the derivative comprises a detection reagent.

5. The method of claims 1 or 3 where the derivative comprises a dye-label.

6. The method of claims 1 or 3 where the derivative comprises a dye selected from the group consisting of TAMRA, ROX, R6G, Fluorescein-12, IRD40, IRD700, BODIPPY®TR, BODIPY®TMR, BODIPY®R6G and BODIPY®FI.

-48-

7. The method of claims 2 or 3 where the acyclonucleotide is radioactively labeled.

5 8. The method of claims 1-3 wherein the DNA polymerase has at least about 20% primary amino acid sequence identity with Vent® DNA polymerase.

10 9. The method of claims 1-3 wherein the DNA polymerase has at least about 30% primary amino acid sequence identity with Vent® DNA polymerase.

15 10. The method of claims 1-3 wherein the DNA polymerase has at least about 70% primary amino acid sequence identity with Vent® DNA polymerase.

11. The method of claims 1-3 wherein the DNA polymerase binds to an antibody probe that has antigenic specificity to Vent® DNA polymerase.

20 12. The method of claims 1-3 wherein the DNA polymerase is encoded by an isolated DNA fragment that hybridizes in a Southern blot to an isolated DNA fragment selected from the group consisting of a DNA fragment having nucleotides 1-1274 of SEQ ID NO:4, a DNA fragment having nucleotides 291-1772 of SEQ ID NO:4, a DNA fragment having
25 nucleotides 3387-3533 of SEQ ID NO:4, a DNA fragment having nucleotides 4704-5396 of SEQ ID NO:4, and a DNA fragment having nucleotides 4718-5437 of SEQ ID NO:4, wherein hybridization is conducted under the following conditions: a) hybridization: 0.75 M NaCl, 0.15 M Tris, 10 mM EDTA, 0.1% sodium pyrophosphate, 0.1% sodium lauryl sulfate,
30 0.03% BSA, 0.03% Ficoll 400, 0.03% PVP and 100 µg/ml boiled calf thymus DNA at 50°C for about 12 hours and; b) wash: 3X30 minutes with 0.1X SET, 0.1% SDS, 0.1% sodium pyrophosphate and 0.1 M phosphate buffer at 45°C.

13. The method of claims 1-3 wherein the DNA polymerase is selected from the group consisting of Vent®, Deep Vent®, *Pfu* and 9°N™ DNA polymerase.

5 14. The method of claims 1-3 wherein the DNA polymerase has been mutated by substitution of the conserved amino acid residue corresponding to Vent® DNA polymerase A488, L492, A493 or Y499.

10 15. The method of claims 1-3 wherein the DNA polymerase has been mutated by substitution of the amino acid residue corresponding to Vent® DNA polymerase A488 by L, I, V, F, S or C.

15 16. The method of claims 1-3 wherein the DNA polymerase has been mutated by substituting the amino acid residue corresponding to Vent® DNA polymerase A488 by L.

20 17. The method of claims 1-3 wherein the DNA polymerase has been mutated by substitution of the amino acid residue corresponding to Vent® DNA polymerase Y499 to L.

25 18. The method of claims 1-3 wherein the DNA polymerase is selected from the group consisting of Vent® (A488L), Vent® (Y499L) and 9°N™ (A485L) DNA polymerases.

30 19. The method of claims 2 or 3 wherein the extent of acyclonucleotide incorporation is greater than that of the corresponding dideoxynucleotide.

35 20. The method of claims 2 or 3 wherein the extent of acyclonucleotide incorporation is at least, approximately, two-fold greater than incorporation of the corresponding dideoxynucleotide.

21. The method of claims 2 or 3 wherein the extent of acyclonucleotide incorporation is at least, approximately, five-fold greater than incorporation of the corresponding dideoxynucleotide.

22. The method of claims 2 or 3 wherein the extent of acyclonucleotide incorporation is at least, approximately, nine-fold greater than incorporation of the corresponding dideoxynucleotide.

23. The method of claims 2 or 3 wherein the extent of ROX-acyclo-CTP incorporation is greater than that of ROX-ddCTP.

24. The method of claims 2 or 3 wherein the extent of ROX-acyclo-CTP incorporation is at least, approximately, two-fold greater than that of ROX-ddCTP.

25. The method of claims 2 or 3 wherein the extent of ROX-acyclo-CTP incorporation is at least, approximately, five-fold greater than that of ROX-ddCTP.

26. The method of claims 2 or 3 wherein the extent of ROX-acyclo-CTP incorporation is at least, approximately, ten-fold greater than that of ROX-ddCTP.

27. The method of claims 1-3 wherein the DNA polymerase is additionally thermostable.


28. The method of claims 1-3 wherein the DNA polymerase has no detectable exonuclease activity.

29. The method of claims 1-3 wherein the DNA polymerase has less than about 5% of the exonuclease activity of the unmodified enzyme.

30. The method of claims 1-3 wherein the DNA polymerase has less than about 25% of the exonuclease activity of the unmodified enzyme.

31. The method of claims 1-3 further comprising the step of employing the resulting sequence-specific termination product or products in DNA sequence determination.

Figure 1A.



2/13

Analog : dNTP




1 : 1 ddCTP



1 : 1 ROX-ddCTP

1 : 10




1 : 1 IRD40-ddCTP

1 : 10



1 : 1 IRD700-ddCTP

1 : 10



1 : 1 Cyanine3-ddCTP

1 : 10



1 : 1 Cyanine5-ddCTP


1 : 10

dNTP



1 : 1 TAMRA-ddCTP

1 : 10




1 : 1 R6G-ddCTP

1 : 10



1 : 1 JOE-ddCTP

1 : 10



1 : 1 R110-ddCTP

1 : 10



1 : 1 FAM-ddCTP

1 : 10



1 : 1 BODIPY® TR-ddCTP

1 : 10



1 : 1 BODIPY® TMR-ddCTP

1 : 10



1 : 1 BODIPY® R6G-ddCTP


1 : 10

1 : 1 BODIPY® FL-ddCTP


1 : 10

Figure 1B.


A.



Analog : dNTP
ddCTP
1 : 1 ROX-ddCTP
1 : 10
1 : 1 ROX-acycloCTP
1 : 10




1 : 1 IR700-ddCTP
1 : 10
1 : 1 IR700-acycloCTP
1 : 10



1 : 1 TAMRA-ddCTP
1 : 10
1 : 1 TAMRA-acycloCTP
1 : 10


B.



Analog : dNTP
ddCTP
1 : 1 ROX-ddCTP
1 : 10
1 : 1 ROX-acycloCTP
1 : 10



1 : 1 IR700-ddCTP
1 : 10
1 : 1 IR700-acycloCTP
1 : 10



1 : 1 TAMRA-ddCTP
1 : 10
1 : 1 TAMRA-acycloCTP
1 : 10

Figure 2.




Figure 3.

Analog : dNTP









Figure 4.



Vent[®] exo- exo-/A488L exo-/Y499L

1 : 1 ddCTP:dNTP				1 : 1 ddCTP:dNTP				1 : 1 ddCTP:dNTP			
ROX-ddCTP:dNTP				ROX-ddCTP:dNTP				ROX-ddCTP:dNTP			
											
1 : 1	1 : 5	1 : 25	dNTP	1 : 1	1 : 5	1 : 25	dNTP	1 : 1	1 : 5	1 : 25	dNTP

primer

Figure 5.




Figure 6.




Figure 7.




Figure 8.




Figure 9.




Figure 10.

A. AmpliTaq® DNA pol, FS

B. 9°N™ exo-/A485L




Figure 10.

A. AmpliTaq® DNA pol, FS

B. 9°N™ exo-/A485L




Figure 10.

A. AmpliTaq® DNA pol, FS

B. 9°N™ exo-/A485L

SEQUENCE LISTING

<110> JACK, WILLIAM E.
GARDNER, ANDREW
NEW ENGLAND BIOLABS, INC.

<120> INCORPORATION OF MODIFIED NUCLEOTIDES BY ARCHAEON DNA
POLYMERASES AND RELATED METHODS

<130> NEB-166-PCT

<140>

<141>

<150> 60/157,204

<151> 1999-09-30

<160> 30

<170> PatentIn Ver. 2.0

<210> 1

<211> 36

<212> DNA

<213> Synthetic Oligonucleotides

<400> 1

caggcagagg cttataaaaa tcctcgccaa cagctt

36

<210> 2

<211> 26

<212> DNA

<213> Synthetic Oligonucleotides

<400> 2

ggtggcagca gccaaactcag cttcct

26

<210> 3

<211> 24

<212> DNA

<213> Synthetic Oligonucleotides

<400> 3

gattctcatg ataagctacg ccga

24

<210> 4

<211> 5837

<212> DNA

<213> *Thermococcus litoralis*

<400> 4

```

gaattcgcga taaaatctat tttcttcttc cattttttcaa tttcaaaaac gtaagcatga 60
gccaaacctc tcgccctttc tctgtccttc ccgctaacc ctttgaaaac tctctccaaa 120
gcattttttg atgaaagctc acgctcctct atgaggggtca gtatatctgc aatgagttcg 180
tgaaggggta ttctgtagaa caactccatg attttcgatt tggatggggg tttaaaaatt 240
tggcggaact tttatttaat ttgaactcca gtttatatct ggtgggtatt atgatactgg 300
acactgatta cataacaaaa gatggcaagc ctataatccg aattttttaag aaagagaacg 360
gggagtttaa aatagaactt gaccctcatt ttcagcccta tatatatgct cttctcaaag 420
atgactccgc tattgaggag ataaaggcaa taaagggcga gagacatgga aaaactgtga 480
gagtgtctga tgcagtgaat gtcaggaaaa aatttttggg aagggaagtt gaagtctgga 540
agctcatttt cgagcatccc caagacgttc cagctatgcg gggcaaaaata agggaacatc 600
cagctgtggt tgacatttac gaatatgaca taccctttgc caagcggtat ctcatagaca 660
agggttgat tcccatggag ggagacgagg agcttaagct ccttgccttt gatattgaaa 720
cgttttatca tgaggagat gaatttgga agggcgagat aataatgatt agttatgccg 780
atgaagaaga ggccagagta atcacatgga aaaatatcga tttgccgtat gtcgatgttg 840
tgtccaatga aagagaaatg ataaagcggt ttgttcaagt tgttaaagaa aaagaccccg 900
atgtgataat aacttacaat ggggacaatt ttgatttgcc gtatctcata aaacgggcag 960
aaaagctggg agttcggctt gtcttaggaa gggacaaaga acatcccgaa cccaagattc 1020
agaggatggg tgatagtttt gctgtggaaa tcaagggtag aatccacttt gatcttttcc 1080
cagttgtgcg aaggacgata aacctcccaa cgtatacgct tgaggcagtt tatgaagcag 1140
ttttaggaaa aacaaaaagc aaattaggag cagaggaaat tgccgctata tgggaaacag 1200
aagaaagcat gaaaaaacta gccagtgact caatggaaga tgctagggca acgtatgagc 1260
tcgggaagga attcttcccc atggaagctg agctggcaaa gctgataggt caaagtgtat 1320
gggacgtctc gagatcaagc accggcaacc tcgtggagtg gtatctttta agggtagcat 1380
acgcgaggaa tgaacttgca ccgaacaaac ctgatgagga agagtataaa cggcgcttaa 1440
gaacaactta cctgggagga tatgtaaaag agccagaaaa aggtttgtgg gaaaatatca 1500
tttatttgga tttccgcagt ctgtaccctt caataatagt tactcacaac gtatccccag 1560
atacccttga aaaagagggc tgtaagaatt acgatgttgc tccgatagta ggatataggt 1620
tctgcaagga ctttccgggc tttattccct ccatactcgg ggacttaatt gcaatgaggc 1680
aagatataaa gaagaaaatg aaatccacaa ttgacccgat cgaaaagaaa atgctcgatt 1740
ataggcaaaag ggctattaaa ttgcttgcaa acagcatctt acccaacgag tggttacca 1800
taattgaaaa tggagaaata aaattcgtga aaattggcga gtttataaac tcttacatgg 1860
aaaaacagaa ggaaaacggt aaaacagtag agaatactga agttctcgaa gtaaacacc 1920
tttttgcatc ctcatccaac aaaaaaatca aagaaagtga agtcaaaaaa gtcaaagccc 1980
tcataagaca taagtataaa gggaaagctt atgagattca gcttagctct ggtagaaaaa 2040
ttaacataac tgctggccat agtctgttta cagttagaaa tggagaaata aaggagttt 2100
ctggagatgg gataaaagaa ggtgacctta ttgtagcacc aaagaaaatt aaactcaatg 2160
aaaaaggggt aagcataaac attccccagt taatctcaga tctttccgag gaagaaacag 2220
ccgacattgt gatgacgatt tcagccaagg gcagaaagaa cttcttttaa ggaatgctga 2280
gaactttaag gtggatgttt ggagaagaaa atagaaggat aagaacattt aatcgctatt 2340
tgttccatct cgaaaaacta ggccttatca aactactgcc ccgcggatat gaagttactg 2400
actgggagag attaaagaaa tataacaac tttacgagaa gcttgctgga agcgtaagt 2460
acaacggaaa caagagagag tatttagtaa tgttcaacga gatcaaggat tttatatctt 2520
acttcccaca aaaagagctc gaagaatgga aaattggaac tctcaatggc tttagaacga 2580
attgtattct caaagtcgat gaggattttg ggaagctcct aggttactat gttagtggag 2640
gctatgcagg tgcacaaaaa aataaaaactg gtggtatcag ttattcgggtg aagctttaca 2700

```

```

atgaggaccc taatgttctt gagagcatga aaaatgtttg agaaaaattc tttggcaagg 2760
ttagagttga cagaaattgc gtaagtatat caaagaagat ggcataacta gttatgaaat 2820
gcctctgtgg agcattagcc gaaaacaaga gaattccttc tgttatactc acctctcccg 2880
aaccggtacg gtggtcattt ttagaggcgt attttacagg cgatggagat atacatccat 2940
caaaaagggt taggctctca acaaaaagcg agctccttgc aaatcagctt gtgttcttgc 3000
tgaactcttt gggaaatatcc tctgtaaaga taggctttga cagtgggggc tatagagtgt 3060
atataaatga agacctgcaa tttccacaaa cgtctaggga gaaaaacaca tactactcta 3120
acttaattcc caaagagatc cttagggacg tgtttggaaa agagttccaa aagaacatga 3180
cgttcaagaa atttaaagag cttgttgact ctggaaaact taacagggag aaagccaagc 3240
tcttgaggtt cttcattaat ggagatatgt tccttgacag agtcaaaagt gttaaagaaa 3300
aggactatga agggtatgtc tatgacctaa gcgttgagga taacgagAAC tttcttggtg 3360
gttttggttt gctctatgct cacaacagct attacggcta tatgggggat cctaaggcaa 3420
gatggctact gaaggaatgt gctgaaagcg ttaccgcatg ggggagacac tacatagaga 3480
tgacgataag agaaatagag gaaaagttcg gctttaaggT tctttatgcg gacagtgtct 3540
caggagaaag tgagatcata ataaggcaaa acggaaagat tagatttgtg aaaataaagg 3600
atcttttctc taaggtggac tacagcattg gcgaaaaaga atactgcatt ctcgaaaggT 3660
ttgaagcact aactctggac gatgacggaa agcttgtctg gaagcccgtc ccctacgtga 3720
tgaggcacag agcgaataaa agaattgttc gcactctggct gaccaacagc tggtatatag 3780
atgttactga ggatcattct ctcataggct atctaaacac gtcaaaaacg aaaactgcca 3840
aaaaaatcgg ggaaagacta aaggaaagtaa agccttttga attaggcaaa gcagtaaaat 3900
cgctcatatg cccaaatgca ccgttaaagg atgagaatac caaaactagc gaaatagcag 3960
taaaattctg ggagctcgta ggattgattg taggagatgg aaactggggT ggagattctc 4020
gttgggcaga gtattatctt ggactttcaa caggcaaaga tgcagaagag ataaagcaaa 4080
aacttctgga acccctaata acttatggag taatctcaaa ctattacca aaaaacgaga 4140
aaggggactt caacatcttg gcaaagagcc ttgtaaagtt tatgaaaagg cactttaagg 4200
acgaaaaagg aagacgaaaa attccagagt tcatgtatga gcttcgggtt acttacatag 4260
aggcatttct acgaggactg ttttcagctg atgggtactgt aactatcagg aagggagttc 4320
cagagatcag gctaacaaac attgatgctg acttttctaag ggaagtaagg aagcttctgt 4380
ggattgttgg aatttcaaat tcaatatttg ctgagactac tccaaatcgc tacaatggtg 4440
tttctactgg aacctactca aagcatctaa ggatcaaaaa taagtggcgt tttgctgaaa 4500
ggataggctt tttaatcgag agaaagcaga agagactttt agaacattta aaatcagcga 4560
gggtaaaaag gaataccata gattttggct ttgatcttgt gcatgtgaaa aaagtcgaag 4620
agataccata cgaggggttac gtttatgaca ttgaagtcca agagacgcag aggttctttg 4680
caaacaacat cctggtacac aatactgacg gcttttatgc cacaataccc ggggaaaagc 4740
ctgaactcat taaaaagaaa gccaaggaat tcctaaacta cataaactcc aaacttccag 4800
gtctgcttga gcttgagtat gagggctttt acttgagagg attctttgtt acaaaaaagc 4860
gctatgcagt catagatgaa gagggcagga taacaacaag gggcttgga gtagtaagga 4920
gagattggag tgagatagct aaggagactc aggcAAaggT tttagaggct atacttaaag 4980
agggaaagtgt tgaAAAagct gtagaagttg ttagagatgt tgtagagaaa atagcaaaat 5040
acagggttcc acttgaaaag cttgttatcc atgagcagat taccagggat ttaaaggact 5100
acaaagccat tggccctcat gtcgcatag caaaaagact tgccgcaaga gggataaaaag 5160
tgaaaccggg cacaataata agctatatcg ttctcaaagg gagcggaaag ataagcgata 5220
gggtaatttt acttacagaa tacgatccta gaaaacacaa gtacgatccg gactactaca 5280
tagaaaacca agttttgccg gcagtactta ggatactcga agcgtttgga tacagaaagg 5340
aggatttaag gtatcaaagc tcaaaacaaa ccggcttaga tgcattggctc aagaggtagc 5400
tctgttgctt tttagtccaa gtttctccgc gagtctctct atctctcttt tgtattctgc 5460
tatgtggttt tcattcacta ttaagtagtc cgccaaagcc ataacgcttc caattccaaa 5520
cttgagctct ttccagctct tggcctcaaa ttcactccat gtttttggat cgtcgtctct 5580

```

```

ccctcttctg ctaagcctct cgaatctttt tcttggcgaa gagtgtacag ctatgatgat 5640
tatctcttcc tctggaaacg catcttttaa cgtctgaatt tcatctagag acctcactcc 5700
gtcgattata actgccttgt acttcttttag tagttctttt acctttggga tcgttaattt 5760
tgccacggca ttgtcccca gctcctgcct aagctgaatg ctcacactgt tcataccttc 5820
gggagttctt gggatcc 5837

```

<210> 5

<211> 15

<212> PRT

<213> *Thermococcus litoralis*

<400> 5

```

Ala Ile Lys Leu Leu Ala Asn Ser Tyr Tyr Gly Tyr Met Gly Tyr
  1             5             10             15

```

<210> 6

<211> 15

<212> PRT

<213> *Pyrococcus* Sp. (GB-D)

<400> 6

```

Ala Ile Lys Ile Leu Ala Asn Ser Tyr Tyr Gly Tyr Tyr Gly Tyr
  1             5             10             15

```

<210> 7

<211> 15

<212> PRT

<213> *Thermococcus* sp.

<400> 7

```

Ala Ile Lys Ile Leu Ala Asn Ser Phe Tyr Gly Tyr Tyr Gly Tyr
  1             5             10             15

```

<210> 8

<211> 15

<212> PRT

<213> *Pyrococcus furiosus*

<400> 8

```

Ala Ile Lys Leu Leu Ala Asn Ser Phe Tyr Gly Tyr Tyr Gly Tyr
  1             5             10             15

```

<210> 9

<211> 15

<212> PRT

<213> *Thermococcus fumicolans*

<400> 9

Ala Ile Lys Ile Leu Ala Asn Ser Phe Tyr Gly Tyr Tyr Gly Tyr
1 5 10 15

<210> 10

<211> 15

<212> PRT

<213> *Thermococcus gorgonarius*

<400> 10

Ala Ile Lys Ile Leu Ala Asn Ser Phe Tyr Gly Tyr Tyr Gly Tyr
1 5 10 15

<210> 11

<211> 15

<212> PRT

<213> *Thermococcus* sp. (TY)

<400> 11

Ala Val Lys Leu Leu Ala Asn Ser Tyr Tyr Gly Tyr Met Gly Tyr
1 5 10 15

<210> 12

<211> 15

<212> PRT

<213> *Pyrococcus abyssi*

<400> 12

Ala Ile Lys Ile Leu Ala Asn Ser Tyr Tyr Gly Tyr Tyr Gly Tyr
1 5 10 15

<210> 13

<211> 15

<212> PRT

<213> *Pyrococcus glycoaerans*

<400> 13

Ala Ile Lys Ile Leu Ala Asn Ser Tyr Tyr Gly Tyr Tyr Gly Tyr
1 5 10 15

<210> 14
<211> 15
<212> PRT
<213> *Pyrococcus horikoshii*

<400> 14
Ala Ile Lys Ile Leu Ala Asn Ser Tyr Tyr Gly Tyr Tyr Gly Tyr
1 5 10 15

<210> 15
<211> 15
<212> PRT
<213> *Pyrococcus* sp. (GE23)

<400> 15
Ala Ile Lys Ile Leu Ala Asn Ser Tyr Tyr Gly Tyr Tyr Gly Tyr
1 5 10 15

<210> 16
<211> 15
<212> PRT
<213> *Pyrococcus* Sp. (KOD1)

<400> 16
Ala Ile Lys Ile Leu Ala Asn Ser Tyr Tyr Gly Tyr Tyr Gly Tyr
1 5 10 15

<210> 17
<211> 15
<212> PRT
<213> *Pyrococcus woesei*

<400> 17
Ala Ile Lys Leu Leu Ala Asn Ser Phe Tyr Gly Tyr Tyr Gly Tyr
1 5 10 15

<210> 18
<211> 15
<212> PRT
<213> *Archaeoglobus fulgidus*

<400> 18
Thr Leu Lys Val Leu Thr Asn Ser Phe Tyr Gly Tyr Met Gly Trp
1 5 10 15

<210> 19

<211> 15

<212> PRT

<213> *Cenarchaeum symbiosum*

<400> 19

Ala	Leu	Lys	Val	Val	Leu	Asn	Ala	Ser	Tyr	Gly	Val	Met	Gly	Ala
1				5					10				15	

<210> 20

<211> 15

<212> PRT

<213> *Methanococcus jannaschii*

<400> 20

Ser	Ile	Lys	Ile	Leu	Ala	Asn	Ser	Val	Tyr	Gly	Tyr	Leu	Ala	Phe
1				5					10				15	

<210> 21

<211> 15

<212> PRT

<213> *Methanococcus voltae*

<400> 21

Ser	Ile	Lys	Val	Leu	Ala	Asn	Ser	His	Tyr	Gly	Tyr	Leu	Ala	Phe
1				5					10				15	

<210> 22

<211> 15

<212> PRT

<213> *Pyrodictium occultum*

<400> 22

Ala	Leu	Lys	Val	Leu	Ala	Asn	Ala	Ser	Tyr	Gly	Tyr	Met	Gly	Trp
1				5					10				15	

<210> 23

<211> 15

<212> PRT

<213> *Sulfurisphaera ohwakuensis*

<400> 23

Ala Met Lys Val Phe Ile Asn Ala Thr Tyr Gly Val Phe Gly Ala
1 5 10 15

<210> 24

<211> 15

<212> PRT

<213> Sulfolobus acidocaldarius

<400> 24

Ala Met Lys Val Phe Ile Asn Ala Thr Tyr Gly Val Phe Gly Ala
1 5 10 15

<210> 25

<211> 15

<212> PRT

<213> Sulfolobus solfataricus

<400> 25

Ala Met Lys Val Phe Ile Asn Ala Thr Tyr Gly Val Phe Gly Ala
1 5 10 15

<210> 26

<211> 15

<212> PRT

<213> Herpesvirus

<400> 26

Ala Ile Lys Val Val Cys Asn Ser Val Tyr Gly Phe Thr Gly Val
1 5 10 15

<210> 27

<211> 15

<212> PRT

<213> human herpesvirus 2

<400> 27

Ala Ile Lys Val Val Cys Asn Ser Val Tyr Gly Phe Thr Gly Val
1 5 10 15

<210> 28

<211> 15

<212> PRT

<213> Human cytomegalovirus

<400> 28

Ala Leu Lys Val Thr Cys Asn Ala Phe Tyr Gly Phe Thr Gly Val
1 5 10 15

<210> 29

<211> 15

<212> PRT

<213> Human DNA Polymerase alpha

<400> 29

Ala Leu Lys Leu Thr Ala Asn Ser Met Tyr Gly Cys Leu Gly Phe
1 5 10 15

<210> 30

<211> 15

<212> PRT

<213> Phage T4

<400> 30

Asn Arg Lys Ile Leu Ile Asn Ser Leu Tyr Gly Ala Leu Gly Asn
1 5 10 15